



BLOW-UP FOR LOGARITHMIC VISCOELASTIC BOUSSINESQ EQUATIONS WITH DELAYED DISSIPATIVE TERM AND ACOUSTIC BOUNDARY CONDITIONS

Amina Benramdane¹, Fatna Bensaber² and Nadia Mezouar³

¹Higher School of Management Tlemcen 13000, Algeria
Statistic and Random Modelization Laboratory, University of Tlemcen, Algeria
e-mail: a.benramdane@esm-tlemcen.dz

²Statistic and Random Modelization Laboratory, University of Tlemcen, Algeria
e-mail: bensaberfatna@yahoo.fr

³Faculty of Economics Sciences, Mascara University, Algeria
e-mail: nadia.mezouar@univ-mascara.dz

Abstract. The current work is devoted to the study of nonlinear viscoelastic wave equations owing a logarithmic source, nonlinear damping and a delay in the velocity with acoustic boundary conditions for which we provide a blow-up criterion. By the use of an energy method, we prove that the solution with negative initial energy blows up after finite time. Moreover, we examine the blow-up time upper and lower bounds.

1. INTRODUCTION

The prediction of waves for coastal regions has always been a big challenge for scientists since wave transformation harms constantly the property and humans near coastal regions. The non-linear transformation of water surface waves in coastal region with the effects of shoaling, diffraction, reflection and refraction are modeled by the use of Boussinesq equations which describe the shallow water waves in intermediate water depth and dissemination of strong

⁰Received July 18, 2025. Revised September 06, 2025. Accepted September 15, 2025.

⁰2020 Mathematics Subject Classification: 35B44, 34M35, 35B05, 58J45.

⁰Keywords: Blow up, local solutions, logarithmic source, nonlinear equation.

⁰Corresponding author: F. Bensaber(bensaberfatna@yahoo.fr).

long waves in surfing areas where breaking of the wave dominates [15]. These equations were formulated by Boussinesq in 1871 and 1872 [3] in order to provide a logical explanation to the existence of the solitary, long, shallow and water waves of permanent form, that was discovered by Scott Russell [16]. The the Boussinesq equations are a system of non linear partial differential equations that consider the influence of wave frequency on the behavior of waves. They are widely applied in numerical description of fairly long waves in the field of coastal engineering to model water waves in shallow seas and ports and weirs [2, 26].

Substantial mathematical interest has been given to Boussinesq equations which was investigated by several researchers from various counts, among them we mention without limitation Valdamir Valmarov [18, 21, 22, 23] and numerous papers such as [4, 9, 24, 25] have investigated the Boussinesq equation generalization. It is important to highlight that considerable amount of effort have been made to establish sufficient conditions for the nonexistence of global solutions to various associated boundary value problems [11, 19]. The effect of small nonlinearity and dispersion in Boussinesq equations is considered, but in many real situations, the damping effect is strong compared with the nonlinear and dispersive one see [8, 21, 22].

Several researcher have studied the blow-up in the Boussinesq equations. For instance, the author in [17] showed the collapse dynamics existence for the Boussinesq equation two basic forms in the case of the periodic boundary conditions. The sufficient criterion of the blow-up is found analytically. Whereas, in [27, 28], the author has studied respectively the existence and the blow-up of solutions to the initial boundary value problems of type Boussinesq equations. An other example ,to name but a few, is in [12] where the author investigated instability of the solitary waves for the generalized Boussinesq equation and established certain blow-up results for improved Boussinesq type equation.

In the current paper, we establish a blow-up result for solutions with vanishing initial energy of the following problem that involves a nonlinear viscoelastic equation of Boussinesq type owing a nonlinear damping term, logarithmic source and a delay in the velocity plus acoustic boundary conditions:

$$\left\{ \begin{array}{ll} u_{tt} + \Delta^2 u - \Delta u + \int_0^t g(t-s)\Delta u(s)ds + a_1 u_t(t)|u_t(t)|^{m-2} \\ + a_2 u_t(t-r)|u_t(t-r)|^{m-2} = a_3 u|u|^{q-2} \ln |u|, & \text{in } \Omega \times (0, T), \\ u(x, t) = 0, & \text{on } \Gamma_0 \times (0, T), \\ \frac{\partial u}{\partial \nu} + \int_0^t g(t-s)\frac{\partial u(s)}{\partial \nu} ds = h_1(x)\mathcal{Z}_t, & \text{on } \Gamma_1 \times (0, T), \\ u_t + h_2(x)\mathcal{Z}_t + h_3(x)\mathcal{Z} = 0, & \text{on } \Gamma_1 \times (0, T), \\ u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x) \text{ in } \Omega, & \mathcal{Z}(x, 0) = \mathcal{Z}_0 \text{ on } \Gamma_1, \\ u_t(x, t-r) = f_0(x, t-r), & \text{in } \Omega \times]0, r[. \end{array} \right. \quad (1.1)$$

With Ω is a bounded domain with a sufficiently smooth boundary in \mathbb{R}^n , $\partial\Omega = \Gamma_1 \cup \Gamma_0$, $\Gamma_1 \cap \Gamma_0 = \emptyset$, $|\Gamma_0| \neq 0$, $|\Gamma_1| \neq 0$, $r > 0$ is a time delay, $m \geq 2$, $q > 2$, $a_1 > 0$, $a_2 \in \mathbb{R}^*$, $a_3 > 0$ and $h_1, h_2, h_3 : \Gamma_1 \rightarrow (0, \infty)$ are essentially bounded functions satisfying

$$0 < h_0 \leq h_3, \text{ for all } x \in \Gamma_1,$$

where h_0 is a constant and the function $g : [0, \infty) \rightarrow [0, \infty)$ is nonincreasing differentiable function.

For that, we split the work into two parts: the second section that provides essential results and hypotheses and a third one where blow-up criteria is proved.

2. PRELIMINARIES

Let $V = \{w \in H^2(\Omega) : w = 0 \text{ on } \Gamma_0\}$ and $\langle \cdot, \cdot \rangle$ denote the scalar product in the space $L^2(\Omega)$. $\|\cdot\|_\tau$ and $\|\cdot\|_{\Gamma_1, \tau}$ represent the norms in the spaces $L^\tau(\Omega)$ and $L^\tau(\Gamma_1)$, respectively. Moreover, $\|\cdot\|_Y$ denotes the norm of space Y . $C > 0$ denotes a generic constant. If there is no ambiguity, we omit the variables t and x . We recall the embedding $H_{\Gamma_0}^2(\Omega) \hookrightarrow L^\tau(\Omega)$, where $\tau \geq 2$ and

$$\begin{cases} \tau < \infty & \text{if } n = 1, 2, 3, 4; \\ \tau \leq \frac{2n}{n-4} & \text{if } n \geq 5. \end{cases} \tag{2.1}$$

Let λ_1 be the first eigenvalue of the spectral Dirichlet problem

$$\Delta^2 u = \lambda_1 u, \text{ in } \Omega, \quad u = \frac{\partial u}{\partial \eta} = 0 \text{ in } \Gamma,$$

$$\|\nabla u\|_2 \leq \frac{1}{\sqrt{\lambda_1}} \|\Delta u\|_2. \tag{2.2}$$

As consequence $V \hookrightarrow L^\tau(\Omega)$, with $\|u\|_\tau \leq \frac{C_n}{\sqrt{\lambda_1}} \|\Delta u\|_2$.

Now, we introduce, as in the work of in Nicaise and Pignotti [14], the new variable

$$y(x, \rho, t) = u_t(x, t - r\rho), \quad x \in \Omega, \quad \rho \in (0, 1), \quad t > 0.$$

Then, we obtain

$$ry_t(x, \rho, t) + y_\rho(x, \rho, t) = 0 \quad \text{in } \Omega \times (0, 1) \times (0, +\infty).$$

Then, the problem (1.1) is equivalent to

$$\left\{ \begin{array}{ll} u_{tt} + \Delta^2 u - \Delta u + \int_0^t g(t-s)\Delta u(s)ds + a_1 u_t(t)|u_t(t)|^{m-2} & \text{in } \Omega \times (0, T), \\ + a_2 y(x, 1, t)|y(x, 1, t)|^{m-2} = a_3 u|u|^{q-2} \ln |u|, & \text{in } \Omega \times]0, 1[\times]0, +\infty[, \\ r y_t(x, \rho, t) + y_\rho(x, \rho, t) = 0, & \text{on } \partial\Omega \times [0, \infty[, \\ u(x, t) = 0, & \text{on } \Gamma_1 \times (0, T), \\ \frac{\partial^2 u}{\partial \nu^2} + \frac{\partial u}{\partial \nu} + \int_0^t g(t-s)\frac{\partial u(s)}{\partial \nu} ds = h_1(x)\mathcal{Z}_t, & \text{on } \Gamma_1 \times (0, T), \\ u_t + h_2(x)\mathcal{Z}_t + h_3(x)\mathcal{Z} = 0, & \text{in } \Omega, \\ u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x), & \text{on } \Gamma_1, \\ \mathcal{Z}(x, 0) = \mathcal{Z}_0, & \text{in } \Omega \times]0, 1[, \\ y(x, \rho, 0) = f_0(x, -\rho r) = y_0(x, \rho), & \end{array} \right. \tag{2.3}$$

To state and prove our result, we set the following assumptions:

- **(A1)** The kernel g verifies $g_l := 1 - \int_0^\infty g(s)ds > 0$.
- **(A2)** The coefficients a_1 and a_2 satisfy

$$a_2 \neq 0 \quad \text{and} \quad -a_1 < a_2 < a_1.$$

- **(A3)** $2 < m < q$.
- **(A4)** Let q satisfy

$$\begin{cases} 2 < q < \infty & \text{if } n = 1, 2, 3, 4; \\ 2 < q < \min \left\{ \frac{2(n-2)}{n-4}, \frac{4+n}{n-4} \right\} & \text{if } n \geq 5. \end{cases}$$

- **(A5)** The kernel function g fulfills

$$\int_0^\infty g(s)ds < \frac{q-2}{q-2+\frac{1}{q}}. \tag{2.4}$$

The following theorem establishes the existence and uniqueness of a local solution of our problem, the proof can be determined by combining the proof given in [6, 13].

Theorem 2.1. *Assume that (A1), (A2) and the exponent q satisfies*

$$\begin{cases} 2 < q < \infty & \text{if } n = 1, 2, 3, 4; \\ 2 < q < \frac{2(n-2)}{n-4} & \text{if } n \geq 5. \end{cases} \tag{2.5}$$

Then, for every $(u_0, u_1, \mathcal{Z}_0, y_0) \in V \times L^2(\Omega) \times L^2(\Gamma) \times L^m(\Omega \times (0, 1))$, problem (2.3) admit a unique local solution (u, \mathcal{Z}, y) with $u \in C(0, T; V) \cap C^1(0, T; L^2(\Omega))$, $\sqrt{h_1}\mathcal{Z} \in L^\infty(0, T; L^2(\Gamma_1))$ and $y \in L^\infty(0, T; L^m(\Omega \times (0, 1)))$.

Our objective is to determine a blow-up criterion to problem (1.1).

3. BLOW-UP CRITERIA

We define the energy associated to the solution of system (1.1) by

$$\begin{aligned}
 E(t) = & \frac{1}{2} \|\Delta u\|_2^2 + \frac{1}{2} \|u_t\|_2^2 + \frac{1}{2} \|\nabla u\|_2^2 - \frac{1}{2} \int_0^t g(s) ds \|\nabla u\|_2^2 \\
 & + \frac{1}{2} (g \circ \nabla u) + \frac{a_3}{q^2} \|u\|_q^q - \frac{a_3}{q} \int_{\Omega} |u|^q \ln |u| dx \\
 & + r \xi \int_0^1 \|y(x, \rho, t)\|_m^m d\rho + \frac{1}{2} \int_{\Gamma_1} h_3 h_1 \mathcal{Z}^2 d\Gamma,
 \end{aligned} \tag{3.1}$$

where

$$(g \circ \nabla u)(t) = \int_0^t g(t-s) \|\nabla u(t) - \nabla u(s)\|_2^2 ds$$

and ξ is a positive constant such that

$$\frac{|a_2|(m-1)}{m} < \xi < \frac{ma_1 - |a_2|}{m} \tag{3.2}$$

and

$$a_1 > |a_2|.$$

Lemma 3.1. *Under the conditions of Theorem 2.1, it holds*

$$\begin{aligned}
 \frac{d}{dt} E(t) \leq & \frac{1}{2} (g' \circ \nabla u - g(t) \|\nabla u(t)\|_2^2) \\
 & - \int_{\Gamma_1} h_2 h_1 \mathcal{Z}_t^2 d\Gamma - \gamma_1 (\|u_t\|_m^m + \|y(x, 1, t)\|_m^m),
 \end{aligned} \tag{3.3}$$

where γ_1 is given later.

Proof. By multiplying the first equation in (2.3) by u_t , integrating over Ω , and using integration by parts, we obtain

$$\begin{aligned}
 & \frac{1}{2} \frac{\partial}{\partial t} \int_{\Omega} |u_t|^2 dx + \int_{\Omega} \Delta u \Delta u_t dx + \int_{\Gamma_1} u_t \frac{\partial \Delta u}{\partial \nu} + \int_{\Omega} \nabla u \nabla u_t - \int_{\Gamma_1} u_t \frac{\partial u}{\partial \nu} d\mu \\
 & - \int_{\Omega} \int_0^t g(t-s) \nabla u(s) \nabla u_t ds dx + a_1 \int_{\Omega} |u_t|^m dx \\
 & + a_2 \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u_t dx - a_3 \int_{\Omega} u |u|^{q-2} \ln |u| u_t dx \\
 & + \int_{\Gamma_1} \int_0^t g(t-s) u_t(t) \frac{\partial u(s)}{\partial \nu} ds d\mu = 0.
 \end{aligned} \tag{3.4}$$

On the other hand,

$$\int_{\Omega} u |u|^{q-2} \ln |u| u_t dx = \frac{d}{dt} \left[\frac{1}{q} \int_{\Omega} |u|^q \ln |u| dx - \frac{1}{q^2} \|u\|_q^q \right], \tag{3.5}$$

$$\begin{aligned} & \int_{\Omega} \int_0^t g(t-s) \nabla u(s) \nabla u_t ds dx \\ &= -\frac{g(t)}{2} \|\nabla u\|_2^2 + \frac{1}{2} (g' \circ \nabla u)(t) + \frac{1}{2} \frac{d}{dt} \left[\int_0^t g(s) ds \|\nabla u\|_2^2 - (g \circ \nabla u)(t) \right]. \end{aligned} \tag{3.6}$$

Then, substituting (3.5) and (3.6) into (3.4), we obtain

$$\begin{aligned} & \frac{1}{2} \frac{\partial}{\partial t} \left[\|u_t\|_2^2 + \|\Delta u\|_2^2 + \|\nabla u\|_2^2 + \int_0^t g(s) ds \|\nabla u\|_2^2 + (g \circ \nabla u)(t) \right. \\ & \quad \left. - \frac{2a_3}{q} \int_{\Omega} |u|^q \ln |u| dx + \frac{2a_3}{q^2} \|u\|_q^q \right] \\ &= \int_{\Gamma_1} u_t \frac{\partial^2 u}{\partial \nu^2} d\Gamma + \int_{\Gamma_1} u_t \frac{\partial u}{\partial \nu} - \frac{g(t)}{2} \|\nabla u\|_2^2 + \frac{1}{2} (g' \circ \nabla u)(t) - a_1 \int_{\Omega} |u_t|^m dx \\ & \quad - a_2 \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u_t dx - \int_{\Gamma_1} \int_0^t g(t-s) u_t(t) \frac{\partial u(s)}{\partial \nu} ds d\mu. \end{aligned} \tag{3.7}$$

We multiply the second equation in (2.3) by $\xi y(x, \rho, t) |y(x, \rho, t)|^{m-2}$, we integrate the result over $\Omega \times (0, 1)$, we obtain

$$\xi r \frac{\partial}{\partial t} \int_0^1 \|y(x, 1, t)\|^m d\rho = -\xi \|y(x, 1, t)\|_m^m + \xi \|u_t\|_m^m. \tag{3.8}$$

We multiply the third equation in (2.3) by u_t , we integrate the result over Γ_1 , we get

$$\int_{\Gamma_1} u_t \frac{\partial^2 u}{\partial \nu^2} d\Gamma + \int_{\Gamma_1} \frac{\partial u}{\partial \nu} u_t d\Gamma + \int_{\Gamma_1} \int_0^t g(t-s) \frac{\partial u(s)}{\partial \nu} ds d\Gamma = \int_{\Gamma_1} h_1(x) \mathcal{Z}_t d\Gamma. \tag{3.9}$$

We multiply the fourth equation in (2.3) by $h_1 \mathcal{Z}_t$, we integrate the result over Γ_1 , we get

$$\int_{\Gamma_1} h_1 \mathcal{Z}_t u_t = - \int_{\Gamma_1} h_1 h_2(x) \mathcal{Z}_t^2 d\Gamma - \frac{1}{2} \int_{\Gamma_1} \frac{\partial}{\partial t} h_1 h_3(x) \mathcal{Z}^2 d\Gamma. \tag{3.10}$$

Next, substituting (3.10) into (3.9), we obtain

$$\begin{aligned} & \int_{\Gamma_1} u_t \frac{\partial^2 u}{\partial \nu^2} d\Gamma + \int_{\Gamma_1} \frac{\partial u}{\partial \nu} u_t d\Gamma + \int_{\Gamma_1} \int_0^t g(t-s) \frac{\partial u(s)}{\partial \nu} ds d\Gamma \\ &= - \int_{\Gamma_1} h_1 h_2(x) \mathcal{Z}_t^2 d\Gamma - \frac{1}{2} \int_{\Gamma_1} \frac{\partial}{\partial t} h_1 h_3(x) \mathcal{Z}^2 d\Gamma. \end{aligned} \tag{3.11}$$

Substituting (3.11) into (3.7), we get

$$\begin{aligned} & \frac{1}{2} \frac{\partial}{\partial t} \left[\|u_t\|_2^2 - \|\Delta u\|_2^2 + \|\nabla u\|_2^2 - \int_0^t g(s) ds \|\nabla u\|_2^2 + (g \circ \nabla u)(t) \right. \\ & \quad \left. - \frac{2a_3}{q} \int_{\Omega} |u|^q \ln |u| dx + \frac{2a_3}{q^2} \|u\|_q^q + \int_{\Gamma_1} h_1 h_3(x) \mathcal{Z}^2 d\Gamma \right] \\ & = -\frac{g(t)}{2} \|\nabla u\|_2^2 + \frac{1}{2} (g' \circ \nabla u)(t) - a_1 \int_{\Omega} |u_t|^m dx \\ & \quad - a_2 \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u_t dx - \int_{\Gamma_1} h_1 h_2(x) \mathcal{Z}_t^2 d\Gamma. \end{aligned} \tag{3.12}$$

By combining (3.12) and (3.8), we obtain

$$\begin{aligned} \frac{\partial}{\partial t} E(t) & = -\frac{g(t)}{2} \|\nabla u\|_2^2 + \frac{1}{2} (g' \circ \nabla u)(t) - (a_1 - \xi) \|u_t\|_m^m - \xi \|y(x, 1, t)\|_m^m \\ & \quad - a_2 \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u_t dx - \int_{\Gamma_1} h_1 h_2(x) \mathcal{Z}_t^2 d\Gamma. \end{aligned} \tag{3.13}$$

By using Young’s inequality, we get

$$-a_2 \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u_t dx \leq |a_2| \frac{(m-1)}{m} \|y(x, 1, t)\|_m^m + \frac{|a_2|}{m} \|u_t\|_m^m, \tag{3.14}$$

then we estimate $E'(t)$

$$\begin{aligned} E'(t) & \leq -\frac{g(t)}{2} \|\nabla u\|_2^2 + \frac{1}{2} (g' \circ \nabla u)(t) - \gamma_1 (\|u_t\|_m^m + \|y(x, 1, t)\|_m^m) \\ & \quad - \int_{\Gamma_1} h_1 h_2(x) \mathcal{Z}_t^2 d\Gamma, \end{aligned} \tag{3.15}$$

where $\gamma_1 = \min\{a_1 - \xi - \frac{a_2}{m}, \frac{\xi m - |a_2|(m-1)}{m}\}$, we choose $a_1 - \xi - \frac{a_2}{m} > 0$ and $\frac{\xi m - |a_2|(m-1)}{m} > 0$

$$\frac{|a_2|(m-1)}{m} < \xi < \frac{ma_1 - a_2}{m}$$

and

$$a_1 > |a_2| > 0.$$

□

Lemma 3.2. Assume (A1) and if $E(0) < 0$, then

$$\int_{\Omega} |u|^q \ln |u| dx \geq 0 \tag{3.16}$$

and

$$\|u_t\|_m^m + \|y(x, 1, t)\|_m^m \leq \frac{I'(t)}{\gamma_1}, \tag{3.17}$$

where $I(t) = -E(t)$.

Proof. Since the decreasing energy, then

$$-\frac{a_3}{p} \int_{\Omega} |u|^p \ln|u| dx < -E(t) < -E(0) < 0, \tag{3.18}$$

it implies that

$$\frac{a_3}{p} \int_{\Omega} |u|^p \ln|u| dx > 0.$$

From (3.15) and g is a nonincreasing, we have

$$\gamma_1 (\|u_t\|_m^m + \|y(x, 1, t)\|_m^m) \leq -E'(t). \tag{3.19}$$

Then we obtain (3.17). □

Lemma 3.3. ([7]) *Let $w \in H_0^2(\Omega)$ and τ assure $\tau \geq 2$ and*

$$\begin{cases} 2 < \tau < \infty & \text{if } n = 1, 2, 3, 4; \\ 2 < \tau \leq \frac{2(n-2)}{n-4} & \text{if } n \geq 5. \end{cases} \tag{3.20}$$

Then, there exist $c > 1$, for all $\frac{2}{\tau} \leq \frac{s}{\tau} \leq 1$, it holds

$$\|w\|_{\tau}^s \leq c(\|w\|_{\tau}^{\tau} + \|\nabla w\|_2^2).$$

Lemma 3.4. ([7]) *Let $\tau \geq 2$. For $w \in H_0^2(\Omega)$ with (3.16), it holds*

$$\|w\|_{\tau}^{\tau} \leq C \left(\|\Delta w\|_2^2 + \int_{\Omega} |w|^{\tau} \ln|w| dx \right).$$

Lemma 3.5. ([7]) *Let $w \in H_0^2(\Omega)$ satisfy (3.16) and τ verify $\tau \geq 2$ and*

$$\begin{cases} 2 < \tau < \infty & \text{if } n = 1, 2, 3, 4; \\ 2 < \tau \leq \frac{2(n-2)}{n-4} & \text{if } n \geq 5. \end{cases}$$

Then, for all $\frac{2}{\tau} \leq \frac{s}{\tau} \leq 1$,

$$\left(\int_{\Omega} |w|^{\tau} \ln|w| dx \right)^{\frac{s}{\tau}} \leq C \left(\|\Delta w\|_2^2 + \int_{\Omega} |w|^{\tau} \ln|w| dx \right).$$

Lemma 3.6. ([7]) *Let $w \in H_0^2(\Omega)$ satisfy (3.16) and τ verify $\tau \geq 2$ and*

$$\begin{cases} 2 < \tau < \infty & \text{if } n = 1, 2, 3, 4; \\ 2 < \tau \leq \frac{2(n-2)}{n-4} & \text{if } n \geq 5. \end{cases}$$

Then there exists a positive constant C such that

$$\|w\|_\tau^\tau \leq C \left(\|\Delta w\|_{\frac{2\tau}{q}}^{\frac{2\tau}{q}} + \left(\int_\Omega |w|^\tau \ln |w| dx \right)^{\frac{\tau}{q}} \right).$$

Theorem 3.7. *Assume that (A1)-(A5) hold and if $E(0) < 0$, the solution to problems (2.3) blows up after a time*

$$T^* \leq \frac{(1 - \beta) (F(0))^{-\frac{\beta}{1-\beta}}}{\beta c_2}$$

with c_2 nonnegative constant, β satisfies

$$\frac{2(q - m)}{q^2(m - 1)} \leq \beta \leq \min \left\{ \frac{q - m}{q(m - 1)}, \frac{q - 2}{2q} \right\} \tag{3.21}$$

and

$$F(0) = I^{1-\beta}(0) + \varepsilon \langle u_0, u_1 \rangle - \frac{\varepsilon}{2} \int_{\Gamma_1} h_1 h_2 \mathcal{Z}_t^2 d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z}_t u_0 d\Gamma > 0.$$

Proof. Let set

$$F(t) = I^{1-\beta}(t) + \varepsilon \langle u, u_t \rangle - \frac{\varepsilon}{2} \int_{\Gamma_1} h_1 h_2 \mathcal{Z}^2 d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z} u d\Gamma, \tag{3.22}$$

where $\varepsilon > 0$. Differentiating with respect to t and using the first equation of (2.3)

$$\begin{aligned} F'(t) = & (1 - \beta) I^{-\beta}(t) I'(t) + \varepsilon \|u_t\|_2^2 - \varepsilon \int_\Omega \Delta^2 u u dx + \varepsilon \int_\Omega \Delta u u dx \\ & - \varepsilon \int_\Omega \int_0^t g(t - s) \Delta u(s) u(t) ds dx - a_1 \varepsilon \int_\Omega u_t(t) |u_t(t)|^{m-2} u(t) dx \\ & - a_2 \varepsilon \int_\Omega y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx + a_3 \varepsilon \int_\Omega |u|^q \ln |u| \\ & - \varepsilon \int_{\Gamma_1} h_1 h_2 \mathcal{Z} \mathcal{Z}_t d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z}_t u d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z} u_t d\Gamma \end{aligned}$$

$$\begin{aligned}
&= (1 - \beta)I^{-\beta}(t)I'(t) + \varepsilon\|u_t\|_2^2 - \varepsilon \int_{\Omega} |\Delta u|^2 dx \\
&\quad + \varepsilon \int_{\Gamma_1} \frac{\partial^2 u}{\partial \nu^2} u d\Gamma - \varepsilon \int_{\Omega} |\nabla u|^2 dx \\
&\quad + \varepsilon \int_{\Gamma_1} \frac{\partial u}{\partial \nu} u d\Gamma - a_1 \varepsilon \int_{\Omega} u_t(t) |u_t(t)|^{m-2} u(t) dx \\
&\quad - a_2 \varepsilon \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx \\
&\quad + a_3 \varepsilon \int_{\Omega} |u|^q \ln |u| + \varepsilon \int_0^t g(s) ds \|\nabla u\|^2 \\
&\quad + \varepsilon \int_0^t g(t-s) \langle \nabla u(t), \nabla u(s) - \nabla u(t) \rangle ds \\
&\quad - \varepsilon \int_{\Gamma_1} g(t-s) \frac{\partial u}{\partial \nu} u ds d\nu \\
&\quad - \varepsilon \int_{\Gamma_1} h_1 h_2 \mathcal{Z} \mathcal{Z}_t d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z}_t u d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z} u_t d\Gamma \\
&= (1 - \beta)I^{-\beta}(t)I'(t) + \varepsilon\|u_t\|_2^2 - \varepsilon\|\Delta u\|_2^2 dx - \varepsilon(1 - \int_0^t g(s) ds) \|\nabla u\|_2^2 \\
&\quad - a_1 \varepsilon \int_{\Omega} u_t(t) |u_t(t)|^{m-2} u(t) dx - a_2 \varepsilon \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx \\
&\quad + a_3 \varepsilon \int_{\Omega} |u|^q \ln |u| + \varepsilon \int_0^t g(t-s) \langle \nabla u(t), \nabla u(s) - \nabla u(t) \rangle ds \\
&\quad + \varepsilon \int_{\Gamma_1} \frac{\partial^2 u}{\partial \nu^2} u d\Gamma + \varepsilon \int_{\Gamma_1} \frac{\partial u}{\partial \nu} u d\Gamma - \varepsilon \int_{\Gamma_1} g(t-s) \frac{\partial u}{\partial \nu} u ds d\nu \\
&\quad - \varepsilon \int_{\Gamma_1} h_1 h_2 \mathcal{Z} \mathcal{Z}_t d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z}_t u d\Gamma - \varepsilon \int_{\Gamma_1} h_1 \mathcal{Z} u_t d\Gamma \\
&\geq (1 - \beta)I^{-\beta}(t)I'(t) + \varepsilon\|u_t\|_2^2 - \varepsilon\|\Delta u\|_2^2 + \varepsilon \int_{\Omega} |u|^q \ln |u| dx \\
&\quad - \varepsilon \left(1 - \int_0^t g(s) ds \right) \|\nabla u\|_2^2 - a_1 \varepsilon \int_{\Omega} u_t(t) |u_t(t)|^{m-2} u(t) dx \\
&\quad - a_2 \varepsilon \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx \\
&\quad + \varepsilon \int_0^t g(t-s) \langle \nabla u(t), \nabla u(s) - \nabla u(t) \rangle ds + \varepsilon \int_{\Gamma_1} h_1 h_3 \mathcal{Z}^2 d\Gamma.
\end{aligned} \tag{3.23}$$

By Young's inequality, we obtain

$$\begin{aligned} \varepsilon \int_0^t g(t-s) \langle \nabla u(t), \nabla u(s) - \nabla u(t) \rangle ds &\geq -\frac{\varepsilon}{4\eta} \|\nabla u\|^2 - 2 \int_0^t g(s) ds \\ &\quad - \varepsilon \eta \int_0^t g(t-s) \|\nabla u(s) - \nabla u(t)\|_2^2 ds \\ &= -\frac{\varepsilon}{4\eta} \|\nabla u\|_2^2 \int_0^t g(s) ds - \varepsilon \eta g \circ \nabla u. \end{aligned}$$

So, we have

$$\begin{aligned} F'(t) &\geq (1-\beta)I^{-\beta}(t)I'(t) + \varepsilon\|u_t\|_2^2 - \varepsilon\|\Delta u\|_2^2 + \varepsilon \int_{\Omega} |u|^q \ln |u| dx \\ &\quad - \varepsilon \left(1 - \int_0^t g(s) ds + \frac{1}{4\eta} \int_0^t g(s) ds \right) \|\nabla u\|_2^2 \\ &\quad - a_1 \varepsilon \int_{\Omega} u_t(t) |u_t(t)|^{m-2} u(t) dx - a_2 \varepsilon \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx \\ &\quad + \varepsilon \int_{\Gamma_1} h_1 h_3 \mathcal{Z}^2 d\Gamma - \varepsilon \eta g \circ \nabla u + \varepsilon \alpha I(t) - \varepsilon \alpha I(t), \end{aligned}$$

where $\alpha = p(1 - \lambda)$ for some $\lambda \in (0, 1)$. Therefore,

$$\begin{aligned} F'(t) &\geq (1-\beta)I^{-\beta}(t)I'(t) + \varepsilon\left(\frac{\alpha}{2} + 1\right)\|u_t\|_2^2 + \varepsilon\left(\frac{\alpha}{2} - 1\right)\|\Delta u\|_2^2 \\ &\quad + \varepsilon \left(1 - \frac{a_3\alpha}{q} \right) \int_{\Omega} |u|^q \ln |u| dx \\ &\quad + \varepsilon \left(\frac{\alpha}{2} - \frac{\alpha}{2} \int_0^t g(s) ds - 1 + \int_0^t g(s) ds - \frac{1}{4\eta} \int_0^t g(s) ds \right) \|\nabla u\|_2^2 \\ &\quad + \varepsilon \left(\frac{\alpha}{2} - \eta \right) g \circ \nabla u - a_1 \varepsilon \int_{\Omega} u_t(t) |u_t(t)|^{m-2} u(t) dx \\ &\quad - a_2 \varepsilon \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx \\ &\quad + \varepsilon r \xi \alpha \int_0^1 \|y(x, \rho, t)\|_m^m d\rho + \varepsilon \left(\frac{\alpha}{2} + 1 \right) \int_{\Gamma_1} h_1 h_3 \mathcal{Z}^2 d\Gamma + \varepsilon \alpha I(t). \end{aligned} \tag{3.24}$$

From **(A2)** and using Young's inequalities with $\frac{m-1}{m} + \frac{1}{m} = 1$, we get

$$\begin{aligned} a_1 \varepsilon \int_{\Omega} u_t(t) |u_t(t)|^{m-2} u(t) dx + a_2 \varepsilon \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx \\ \leq \frac{a_1(m-1)}{m} \delta_1^{-\frac{m}{m-1}} (\|y(1, t)\|_m^m + \|u_t\|_m^m) + \frac{(a_1 + |a_2|)\delta_1^m}{m} \|u\|_m^m. \end{aligned} \tag{3.25}$$

From (3.17) and taking $\delta_1 = (\theta I^{-\beta}(t))^{-\frac{m-1}{m}}$ and $\theta > 0$, we obtain

$$\begin{aligned} & \frac{a_1(m-1)}{m} \delta_1^{-\frac{m}{m-1}} (\|y(1, t)\|_m^m + \|u_t\|_m^m) + \frac{(a_1 + |a_2|)\delta_1^m}{m} \|u\|_m^m \\ & \leq \frac{a_1(m-1)\theta}{m\gamma_1} (I(t))^{-\beta} I'(t) + \frac{C}{m\theta^{m-1}} (I(t))^{\beta(m-1)} \|u\|_m^m \tag{3.26} \\ & \leq \frac{a_1(m-1)\theta}{m\gamma_1} (I(t))^{-\beta} I'(t) + \frac{C}{m\theta^{m-1}} (I(t))^{\beta(m-1)} \|u\|_q^m. \end{aligned}$$

Through the utility of **(A3)** and **(A4)**. On account of (3.18) and Lemma 3.4, we obtain

$$(I(t))^{\beta(m-1)} \|u\|_q^m \leq C \left(\|\Delta u\|_2^2 + \int_{\Omega} |u|^q \ln|u| dx \right)^{\frac{m}{q}} \left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\beta(m-1)}. \tag{3.27}$$

By the elementary inequality $(a + b)^{\Upsilon} \leq 2^{\Upsilon}(a^{\Upsilon} + b^{\Upsilon})$, we get

$$\begin{aligned} & (I(t))^{\beta(m-1)} \|u\|_q^m \\ & \leq 2^{\frac{m}{q}} C \left(\|\Delta u\|_2^{\frac{2m}{q}} + \left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\frac{m}{q}} \right) \left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\beta(m-1)} \\ & \leq C \left[\|\Delta u\|_2^{\frac{2m}{q}} \left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\beta(m-1)} + \left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\beta(m-1) + \frac{m}{q}} \right]. \tag{3.28} \end{aligned}$$

Using Young’s equality with $\frac{m}{q} + \frac{q-m}{q} = 1$, this estimation becomes

$$\begin{aligned} & (I(t))^{\beta(m-1)} \|u\|_q^m \\ & \leq C \left[\|\Delta u\|_2^2 + \left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\frac{\beta(m-1)q}{q-m}} + \left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\frac{\beta q(m-1)+m}{q}} \right]. \tag{3.29} \end{aligned}$$

By (3.21), we observe that $\frac{2}{q} \leq \frac{\beta q(m-1)+m}{q} \leq 1$ and $\frac{2}{q} \leq \frac{\beta(m-1)q}{q-m} < 1$. Then, utilizing Lemma 3.5, we deduce

$$\left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\frac{\beta q(m-1)+m}{q}} \leq C \left(\|\Delta u\|_2^2 + \int_{\Omega} |u|^q \ln|u| dx \right) \tag{3.30}$$

and

$$\left(\int_{\Omega} |u|^q \ln|u| dx \right)^{\frac{\beta(m-1)q}{q-m}} \leq C \left(\|\Delta u\|_2^2 + \int_{\Omega} |u|^q \ln|u| dx \right). \tag{3.31}$$

By inserting (3.30) and (3.31) to (3.29), we obtain

$$(I(t))^{\beta(m-1)} \|u\|_q^m \leq \tilde{C} \left(\|\Delta u\|_2^2 + \int_{\Omega} |u|^q \ln |u| dx \right). \tag{3.32}$$

Combining (3.32) in (3.26), we obtain

$$\begin{aligned} & a_1 \varepsilon \int_{\Omega} u_t(t) |u_t(t)|^{m-2} u(t) dx + a_2 \varepsilon \int_{\Omega} y(x, 1, t) |y(x, 1, t)|^{m-2} u(t) dx \\ & \leq \frac{a_1(m-1)\theta}{m\gamma_1} (I(t))^{-\beta} I'(t) + \frac{C}{m\theta^{m-1}} \left(\|\Delta u\|_2^2 + \int_{\Omega} |u|^q \ln |u| dx \right). \end{aligned} \tag{3.33}$$

By applying this to (3.24), we arrive at

$$\begin{aligned} F'(t) & \geq \left(1 - \beta - \varepsilon \frac{a_1(m-1)\theta}{m\gamma_1} \right) I^{-\beta}(t) I'(t) + \varepsilon \left(\frac{\alpha}{2} + 1 \right) \|u_t\|_2^2 \\ & + \varepsilon \left(-\frac{C}{m\theta^{m-1}} + \frac{\alpha}{2} - 1 \right) \|\Delta u\|_2^2 + \varepsilon \left(1 - \frac{a_3\alpha}{q} - \frac{c}{m\theta^{m-1}} \right) \int_{\Omega} |u|^q \ln |u| dx \\ & + \varepsilon \left(\frac{\alpha}{2} - 1 - \left(\frac{\alpha}{2} - 1 + \frac{1}{4\eta} \right) \int_0^t g(s) ds \right) \|\nabla u\|_2^2 + \varepsilon \left(\frac{\alpha}{2} - \eta \right) g \circ \nabla u \\ & + \varepsilon \frac{a_3}{p^2} \alpha \|u\|_m^m + \varepsilon \left(\frac{\alpha}{2} + 1 \right) \int_{\Gamma_1} h_1 h_3 \mathcal{Z}^2 d\Gamma + \varepsilon \alpha I(t). \end{aligned}$$

Let take $\eta = \frac{\alpha}{2}$ and use the relation

$$\begin{aligned} & \left(\frac{\alpha}{2} - 1 \right) - \left(\frac{\alpha}{2} - 1 + \frac{1}{4\eta} \right) \int_0^t g(s) ds \\ & = \left(\frac{\alpha}{2} - 1 \right) \left(1 - \int_0^t g(s) ds \right) - \frac{1}{2\alpha} \int_0^t g(s) ds \\ & \geq \left(\frac{\alpha}{2} - 1 \right) g_t - \frac{1}{2\alpha} (1 - g_t) =: a_{\alpha}. \end{aligned}$$

Then from (3.24), we get

$$\begin{aligned} F'(t) & \geq \left(1 - \beta - \varepsilon \frac{a_1(m-1)\theta}{m\gamma_1} \right) I^{-\beta}(t) I'(t) + \varepsilon \left(\frac{\alpha}{2} + 1 \right) \|u_t\|_2^2 \\ & + \varepsilon \left(1 - \frac{a_3\alpha}{q} - \frac{c}{m\theta^{m-1}} \right) \int_{\Omega} |u|^q \ln |u| dx \\ & + \varepsilon a_{\alpha} \|\nabla u\|_2^2 + \left(\frac{\alpha}{2} - 1 - \frac{c}{m\theta^{m-1}} \right) \|\Delta u\|_2^2 \\ & + \varepsilon \frac{a_3}{p^2} \alpha \|u\|_q^q + \varepsilon \left(\frac{\alpha}{2} + 1 \right) \int_{\Gamma_1} h_1 h_3 \mathcal{Z}^2 d\Gamma + \varepsilon \alpha I(t). \end{aligned} \tag{3.34}$$

So,

$$F'(t) \geq \tilde{C} (I(t) + \|u_t\|_2^2 + \|\nabla u\|_2^2 + \|u\|_q^q) > 0. \tag{3.35}$$

Consequently, we have $F(t) \geq F(0) > 0$ for all $t > 0$.

On the one hand, from (3.22), we know that

$$F(t) \leq I^{1-\beta}(t) + \varepsilon \langle u, u_t \rangle,$$

then

$$(F(t))^{\frac{1}{1-\beta}} \leq \left(I^{1-\beta}(t) + \varepsilon \langle u, u_t \rangle \right)^{\frac{1}{1-\beta}}. \tag{3.36}$$

We use Hölder’s and Young’s inequalities to get

$$\begin{aligned} \left| \int_{\Omega} uu_t dx \right|^{\frac{1}{1-\beta}} &\leq C \|u\|_q^{\frac{1}{1-\beta}} \|u_t\|_2^{\frac{1}{1-\beta}} \\ &\leq C \|u\|_q^{\frac{2}{1-2\beta}} + \|u_t\|_2^2 \end{aligned} \tag{3.37}$$

with $2 \leq \frac{2}{1-2\beta} \leq q$.

By using Lemma 3.3, we obtain

$$\|u\|_q^{\frac{2}{1-2\beta}} \leq c (\|u\|_q^q + \|\nabla u\|_2^2). \tag{3.38}$$

Substituting (3.38) into (3.37), we get

$$\left| \int_{\Omega} uu_t dx \right|^{\frac{1}{1-\beta}} \leq C (\|u\|_q^q + \|u_t\|_2^2 + \|\nabla u\|_2^2). \tag{3.39}$$

From (3.36) and (3.39), we have

$$(F(t))^{\frac{1}{1-\beta}} \leq \tilde{C} (I + \|u\|_q^q + \|u_t\|_2^2 + \|\nabla u\|_2^2). \tag{3.40}$$

Combining (3.40) and (3.35), we obtain

$$F'(t) \geq C_2 (F(t))^{\frac{1}{1-\beta}}, \tag{3.41}$$

where C_2 is positive constant.

By integrating of (3.41) over $(0, t)$, we have

$$(F(t))^{\frac{\beta}{1-\beta}} \geq \frac{1}{(F(0))^{\frac{-\beta}{1-\beta}} - C_2 \frac{\beta t}{1-\beta}},$$

which gives that the solution u blow up after finite time

$$T \leq T^* = \frac{(1 - \beta) (F(0))^{\frac{-\beta}{1-\beta}}}{\beta C_2}.$$

□

REFERENCES

- [1] K. Agre and M.A. Rammaha, *Systems of nonlinear wave equations with damping and source terms*, *Diff. Integ. Equ.*, **19**(11) (2006), 1235–1270.
- [2] A.S. Ahmadin, *Numerical Models for submerged breakwaters, Coastal Hydrodynamics and Morphodynamics*, Elsevier, 2016.
- [3] J. Boussinesq, *Théorie des ondes et des remous qui se propagent le long d'un canal rectangulaire horizontal, en communiquant au liquide contenu dans ce canal des vitesses sensiblement pareilles de la surface au fond*, *J. Math. Pures Appl.*, 2e série, **17** (1872), 55–108.
- [4] X. Du, *Classification of Single Traveling Wave Solutions to the Generalized Strong Nonlinear Boussinesq Equation without Dissipation Terms in $P = 1$* , *J. Appl. Math. Phys.*, **2**(3) (2014), 50–59.
- [5] I. Ekeland, *On the variational principle*, *J. Math. Anal. Appl.*, **47** (1974), 324–353.
- [6] L. Gongwei, *The existence, general decay and blow-up for a plate equation with nonlinear damping and a logarithmic source term*, *Elec. Resea. Archive*, **28**(1) (2020), 263–289.
- [7] M. Kafini and S. Messaoudi, *Local existence and blow up of solutions to a logarithmic nonlinear wave equation with delay*, *Appl. Anal.*, **99** (2020), 530–547.
- [8] S. Lai and Y. Wu, *The Asymptotic Solution of the Cauchy Problem for a Generalized Boussinesq Equation*, *Disc. Conti. Dyna. Systems-Series B.*, **3**(3) (2003), 401–408.
- [9] H.A. Levine, *Instability and Nonexistence of Global Solutions to Nonlinear Wave Equations of the Form $Pu_{tt} = -Au + F(u)$* , *Tran. Amer. Math. Soc.*, **192** (1974), 1–21.
- [10] H.A. Levine, *A Note on the Non-existence of Global Solutions of Initial Boundary Value Problems for the Boussinesq Equation $u_{tt} = 3u_{xxxx} - 12(u^2)_{xx}$* , *J. Math. Anal. Appl.*, **107** (1985), 206–210.
- [11] F. Linares and M. Scialom, *Asymptotic behavior of solutions of a generalized Boussinesq type equation*, *Nonlinear Anal.*, **25**(11) (1995), 1147–1158.
- [12] Y. Liu, *Instability of solitary waves for generalized Boussinesq equations*, *J. Dyna. Diff. Equ.*, **5** (1993), 537–558.
- [13] N. Mezouar, M. Abdelli and A. Rachah, *Existence of global solutions and decay estimates for a viscoelastic Petrovsky equation with a delay term in the non-linear internal feedback*, *Elect. J. Diff. Equ.*, **2017**(58) (2017), 1–25.
- [14] S. Nicaise and C. Pignotti, *Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks*, *SIAM J. Cont. Optim.*, **45**(5) (2006), 1561–1585.
- [15] P. Patel, P. Kumar and R. Rajni, *The numerical solution of Boussinesq equation for shallow water waves*, *AIP Conference Proc. AIP Publishing*, **2214**(1) (2020), 1–6.
- [16] J. Scott Russell, *Report on waves*, Report of the 14th Meeting of the British Association for the Advancement of Science, John Murray, London, 1844, 311–390.
- [17] S.K. Turitsyn, *Blow-up in the Boussinesq equation*, *Phys. Rev. E. Amer. Phys. Soc.*, **47**(2) (1993), 796–799.
- [18] V. Varlamov, *On the Cauchy problem for the damped Boussinesq equation*, *Diff. Integ. Equ.*, **9**(3) (1996), 619–634.
- [19] V. Varlamov, *Existence and uniqueness of a solution to the Cauchy problem for the damped Boussinesq equations*, *Math. Meth. Appl. Sci.*, **19**(8) (1996), 639–649.
- [20] V. Varlamov, *Asymptotic behavior of solutions of the damped Boussinesq equation in two space dimensions*, *Int. J. Math. Math. Sci.*, **22** (1999), 131–145.
- [21] V. Varlamov, *On the damped Boussinesq equation in a circle*, *Nonlinear Anal.*, **38**(4) (1999), 447–470.

- [22] V. Varlamov, *On the spatially two-dimensional Boussinesq equation in a circular domain*, *Nonlinear Anal.*, **46**(5) (2001), 699–725.
- [23] V. Varlamov, *Two-dimensional Boussinesq equation in a disc and anisotropic Sobolev spaces*, *Comptes Rendus Mécanique*, **335**(9-10) (2007), 548–558.
- [24] Q. Wang, *Numerical Solutions of the Improved Boussinesq Equation with Stokes Damping*, *Appl. Math.*, **12**(4) (2021), 241–251.
- [25] J.H. Wu, *The 2D Incompressible Boussinesq Equations*, Peking University Summer School Lecture Notes, Beijing, 2012.
- [26] Y.T. Zerihun and J.D. Fenton, *Application of a Boussinesq-type equation to flow over trapezoidal profile weirs*, *Hydraulics of Dams and River Structures*, (2004), 369–376.
- [27] Y. Zhijian, *On local existence of solutions of initial boundary value problems for the bad Boussinesq-type equation*, *Nonlinear Anal.*, **51**(7) (2002), 1259–1271.
- [28] Y. Zhijian and X. Wang, *Blow-up of solutions for improved Boussinesq type equation*, *J. Math. Anal. Appl.*, **278**(2) (2003), 335–353.