



## MAJORIZATION PROBLEMS FOR UNIFORMLY STARLIKE FUNCTIONS BASED ON RUSCHEWEYH $q$ -DIFFERENTIAL OPERATOR RELATED WITH EXPONENTIAL FUNCTION

K. Vijaya<sup>1</sup>, G. Murugusundaramoorthy<sup>2</sup> and N. E. Cho<sup>3</sup>

<sup>1</sup>School of Advanced Sciences  
Vellore Institute of Technology, Vellore - 632014, India  
e-mail: [kvijaya@vit.ac.in](mailto:kvijaya@vit.ac.in)

<sup>2</sup>School of Advanced Sciences  
Vellore Institute of Technology, Vellore - 632014, India  
e-mail: [gmsmoorthy@yahoo.com](mailto:gmsmoorthy@yahoo.com)

<sup>3</sup>Department of Applied Mathematics  
Pukyong National University, Busan 48513, Korea  
e-mail: [necho@pknu.ac.kr](mailto:necho@pknu.ac.kr)

**Abstract.** The main object of this present paper is to study some majorization problems for certain classes of analytic functions defined by means of  $q$ -calculus operator associated with exponential function.

### 1. INTRODUCTION

Let  $\mathcal{A}$  be the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

---

<sup>0</sup>Received July 8, 2020. Revised August 31, 2020. Accepted October 3, 2020.

<sup>0</sup>2010 Mathematics Subject Classification: 30C45, 30C80, 33C10.

<sup>0</sup>Keywords: Analytic functions, starlike and convex functions of complex order, uniformly starlike functions, quasi-subordination, majorization problems, Hadamard product (convolution),  $q$ -calculus operator, exponential function.

<sup>0</sup>Corresponding author: N. E. Cho([necho@pknu.ac.kr](mailto:necho@pknu.ac.kr)).

which are analytic in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ . For given  $g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in \mathcal{A}$ , the Hadamard product of  $f$  and  $g$  is defined by

$$(f * g)(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n = (g * f)(z).$$

For two analytic functions  $f, g \in \mathcal{A}$ , we say that  $f$  is subordinate to  $g$ , denoted by  $f \prec g$ , if there exists a Schwarz function  $\omega(z)$  which is analytic in  $\mathbb{U}$  with  $\omega(0) = 0$  and  $|\omega(z)| < 1$  for all  $z \in \mathbb{U}$ , such that  $f(z) = g(\omega(z))$  for  $z \in \mathbb{U}$ . Note that, if the function  $g$  is univalent in  $\mathbb{U}$ , due to Miller and Mocanu [6], we have

$$f(z) \prec g(z) \iff f(0) = g(0) \text{ and } f(\mathbb{U}) \subset g(\mathbb{U}).$$

If  $f$  and  $g$  are analytic functions in  $\mathbb{U}$ , following MacGregor [5], we say that  $f$  is majorized by  $g$  in  $\mathbb{U}$ , that is  $f(z) \ll g(z)$  ( $z \in \mathbb{U}$ ) if there exists a function  $\phi(z)$ , analytic in  $\mathbb{U}$ , such that

$$|\phi(z)| < 1 \text{ and } f(z) = \phi(z)g(z) \quad (z \in \mathbb{U}).$$

It is of interest to note that the notion of majorization is closely related to the concept of quasi-subordination between analytic functions.

Now we recall here the notion of  $q$ -operator that is,  $q$ -difference operator that play vital role in the theory of hypergeometric series, quantum physics and in the operator theory. The application of  $q$ -calculus was initiated by Jackson [3], recently Kanas and Răducanu [4] have used the fractional  $q$ -calculus operators in investigations of certain classes of functions which are analytic in  $\mathbb{U}$ .

Let  $0 < q < 1$ . For any non-negative integer  $n$ , the  $q$ -integer number  $n$  is defined by

$$[n]_q = \frac{1 - q^n}{1 - q} = 1 + q + \cdots + q^{n-1}, \quad [0]_q = 0. \quad (1.2)$$

In general, we will denote

$$[x]_q = \frac{1 - q^x}{1 - q}$$

for a non-integer number  $x$ . Also the  $q$ -number shifted factorial is defined by

$$[n]_q! = [n]_q [n-1]_q \cdots [2]_q [1]_q, \quad [0]_q! = 1. \quad (1.3)$$

Clearly,

$$\lim_{q \rightarrow 1^-} [n]_q = n \quad \text{and} \quad \lim_{q \rightarrow 1^-} [n]_q! = n!.$$

For  $0 < q < 1$ , the Jackson's  $q$ -derivative operator (or  $q$ -difference operator) of a function  $f \in \mathcal{A}$  given by (1.1) defined as follows [3]:

$$\mathfrak{D}_q f(z) = \begin{cases} \frac{f(z) - f(qz)}{(1-q)z} & \text{for } z \neq 0 \\ f'(0) & \text{for } z = 0 \end{cases}, \quad (1.4)$$

$\mathfrak{D}_q^0 f(z) = f(z)$ , and  $\mathfrak{D}_q^m f(z) = \mathfrak{D}_q(\mathfrak{D}_q^{m-1} f(z))$ ,  $m \in \mathbb{N} = \{1, 2, \dots\}$ . From (1.4), we have

$$\mathfrak{D}_q f(z) = 1 + \sum_{n=2}^{\infty} [n]_q a_n z^{n-1} \quad (z \in \mathbb{U}), \quad (1.5)$$

where  $[n]_q$  is given by (1.2).

For a function  $\psi(z) = z^n$ , we obtain

$$\mathfrak{D}_q \psi(z) = \mathfrak{D}_q z^n = \frac{1 - q^n}{1 - q} z^{n-1} = [n]_q z^{n-1}$$

and

$$\lim_{q \rightarrow 1^-} \mathfrak{D}_q \psi(z) = \lim_{q \rightarrow 1^-} ([n]_q z^{n-1}) = n z^{n-1} = \psi'(z),$$

where  $\psi'$  is the ordinary derivative.

Let  $t \in \mathbb{R}$  and  $n \in \mathbb{N}$ . The  $q$ -generalized Pochhammer symbol is defined by

$$[t; n]_q = [t]_q [t+1]_q [t+2]_q \dots [t+n-1]_q \quad (1.6)$$

and for  $t > 0$  the  $q$ -gamma function is defined by

$$\Gamma_q(t+1) = [t]_q \Gamma_q(t) \quad \text{and} \quad \Gamma_q(1) = 1. \quad (1.7)$$

Using the  $q$ -difference operator, Kannas and Raducanu [4] defined the Ruscheweyh  $q$ -differential operator as below: For  $f \in \mathcal{A}$ ,

$$\mathcal{R}_q^\delta f(z) = f(z) * F_{q, \delta+1}(z) \quad (\delta > -1, z \in \mathbb{U}), \quad (1.8)$$

where

$$F_{q, \delta+1}(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma_q(n+\delta)}{[n-1]_q! \Gamma_q(1+\delta)} z^n = z + \sum_{n=2}^{\infty} \frac{[\delta+1; n-1]_q}{[n-1]_q!} z^n. \quad (1.9)$$

Making use of (1.8) and (1.9), we have

$$\mathcal{R}_q^\delta f(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma_q(n+\delta)}{[n-1]_q! \Gamma_q(1+\delta)} a_n z^n \quad (z \in \mathbb{U}). \quad (1.10)$$

$$\begin{aligned}\mathcal{R}_q^0 f(z) &= f(z), \\ \mathcal{R}_q^1 f(z) &= z\mathfrak{D}_q f(z), \\ \mathcal{R}_q^m f(z) &= \frac{z\mathfrak{D}_q^m(z^{m-1}f(z))}{[m]_q!} \quad (m \in \mathbb{N}).\end{aligned}$$

Also we have

$$\mathfrak{D}_q(\mathcal{R}_q^\delta f(z)) = 1 + \sum_{n=2}^{\infty} \Theta_n(q, \delta) a_n z^{n-1}, \quad (1.11)$$

where

$$\Theta_n := \Theta_n(q, \delta) = \frac{[n]_q \Gamma_q(n + \delta)}{[n-1]_q! \Gamma_q(1 + \delta)}. \quad (1.12)$$

It is easy to check that

$$z\mathfrak{D}_q(F_{q, \delta+1}(z)) = \left(1 + \frac{[\delta]_q}{q^\delta}\right) F_{q, \delta+2}(z) - \frac{[\delta]_q}{q^\delta} F_{q, \delta+1}(z) \quad (z \in \mathbb{U}). \quad (1.13)$$

Making use of (1.8)-(1.13) and the properties of Hadamard product, we obtain the following equality

$$z\mathfrak{D}_q(\mathcal{R}_q^\delta f(z)) = \left(1 + \frac{[\delta]_q}{q^\delta}\right) \mathcal{R}_q^{1+\delta} f(z) - \frac{[\delta]_q}{q^\delta} \mathcal{R}_q^\delta f(z) \quad (z \in \mathbb{U}). \quad (1.14)$$

From (1.10), we note that

$$\lim_{q \rightarrow 1^-} F_{q, \delta+1}(z) = \frac{z}{(1-z)^{\delta+1}}, \quad \lim_{q \rightarrow 1^-} \mathcal{R}_q^\delta f(z) = f(z) * \frac{z}{(1-z)^{\delta+1}}.$$

Thus, when  $q \rightarrow 1^-$  we can say that Ruscheweyh  $q$ -differential operator reduces to the differential operator defined by Ruscheweyh [9] and (1.14) gives the well-known recurrent formula for Ruscheweyh differential operator.

Majorization problems for the class  $\mathcal{S}^* = \mathcal{S}^*(0)$  had been investigated by MacGregor [5], further Altintas et al. [1] investigated a majorization problem for  $\mathcal{S}(\gamma)$  the class of starlike functions of complex order  $\gamma$  ( $\gamma \in \mathbb{C} \setminus \{0\}$ ), and Goyal and Goswami [2] generalized these results for the class of analytic functions involving fractional operator. Very lately, Tang and Deng [12] considered majorization properties for multivalent analytic functions related to the Srivastava-Khairnar-More operator and exponential function.

In this paper, using Ruscheweyh  $q$ -differential operator defined by (1.10) and motivated by recent works of [8], we define a new subclass of uniformly starlike functions associated with  $q$ -calculus operator, which are subordinate to exponential function, and investigate a majorization problem. Further we point out some special cases of our result.

**Definition 1.1.** A function  $f \in \mathcal{A}$  is said to be in the class  $\mathcal{RS}_q^\delta(\beta, e^z)$ , if and only if

$$\left[ \frac{z\mathcal{D}_q(\mathcal{R}_q^\delta f(z))}{\mathcal{R}_q^\delta f(z)} - \beta \left| \frac{z\mathcal{D}_q(\mathcal{R}_q^\delta f(z))}{\mathcal{R}_q^\delta f(z)} - 1 \right| \right] \prec e^z, \quad (1.15)$$

where  $\delta > -1$ ,  $\beta > 0$  and  $z \in \mathbb{U}$ .

For  $\beta = 0$  we have  $\mathcal{RS}_q^\delta(\beta, e^z) \equiv \mathcal{RS}_q^\delta(e^z)$ :

$$\frac{z\mathcal{D}_q(\mathcal{R}_q^\delta f(z))}{\mathcal{R}_q^\delta f(z)} \prec e^z$$

where  $\delta > -1$ ,  $\beta > 0$  and  $z \in \mathbb{U}$ .

Further by taking  $q \rightarrow 1^-$  and  $\delta = 0$  we have  $\mathcal{RS}_q^\delta(e^z) \equiv \mathcal{S}^*(e^z)$ :

$$\frac{zf'(z)}{f(z)} \prec e^z \quad (z \in \mathbb{U}).$$

## 2. A MAJORIZATION PROBLEM FOR THE CLASS $\mathcal{RS}_q^\delta(\beta, e^z)$

We state the following  $q$ -analogue of the result given by Nehari (cf. [7]) and Selvakumaran et al. [10].

**Lemma 2.1.** *If the function  $\phi(z)$  is analytic and  $|\phi(z)| < 1$  in  $\mathbb{U}$ , then*

$$|\mathcal{D}_q\phi(z)| \leq \frac{1 - |\phi(z)|^2}{1 - |z|^2}. \quad (2.1)$$

**Theorem 2.2.** *Let the function  $f \in \mathcal{A}$  and suppose that  $g \in \mathcal{RS}_q^\delta(\beta, e^z)$ . If  $\mathcal{R}_q^\delta f(z)$  is majorized by  $\mathcal{R}_q^\delta g$  in  $\mathbb{U}$ , then*

$$|\mathcal{R}_q^{\delta+1} f(z)| \leq |\mathcal{R}_q^{\delta+1} g(z)| \quad (|z| \leq r_1), \quad (2.2)$$

where  $r_1 = r_1(\delta, \beta)$ , is the smallest positive root of the equation

$$r^2 q^\delta e^r - r^2 \{[\delta]_q - ([\delta]_q + q^\delta)\beta\} - q^\delta e^r - 2rq^\delta(1+\beta) + \{[\delta]_q - ([\delta]_q + q^\delta)\beta\} = 0, \quad (2.3)$$

where  $[\delta]_q > ([\delta]_q + q^\delta)\beta + q^\delta e$  and  $\beta \geq 0$ .

*Proof.* Since  $g \in \mathcal{RS}_q^\delta(\beta, e^z)$ , we find from (1.15) that

$$\left[ \left( \frac{z\mathcal{D}_q(\mathcal{R}_q^\delta g(z))}{\mathcal{R}_q^\delta g(z)} \right) - \beta \left| \frac{z\mathcal{D}_q(\mathcal{R}_q^\delta g(z))}{\mathcal{R}_q^\delta g(z)} - 1 \right| \right] = e^{w(z)}, \quad (2.4)$$

where  $w(z) = c_1z + c_2z^2 + c_3z^3 + \dots$  is analytic in  $\mathbb{U}$ , with  $w(0) = 0$  and  $|w(z)| \leq |z|$  for all  $z \in \mathbb{U}$ . Letting

$$\varpi = \frac{z\mathfrak{D}_q(\mathcal{R}_q^\delta g(z))}{\mathcal{R}_q^\delta g(z)}$$

in (2.4), we have

$$\varpi - \beta|\varpi - 1| = e^{w(z)}$$

which implies

$$\varpi = \frac{e^{w(z)} - \beta e^{-i\varphi}}{1 - \beta e^{-i\varphi}}.$$

This is, from (2.4), we get

$$\frac{z\mathfrak{D}_q(\mathcal{R}_q^\delta g(z))}{\mathcal{R}_q^\delta g(z)} = \frac{e^{w(z)} - \beta e^{-i\varphi}}{1 - \beta e^{-i\varphi}}. \quad (2.5)$$

Now, by applying the relation (1.14) in (2.5), we get

$$\frac{\mathcal{R}_q^{\delta+1}g(z)}{\mathcal{R}_q^\delta g(z)} = \frac{[\delta]_q - ([\delta]_q + q^\delta)\beta e^{-i\varphi} + q^\delta e^{w(z)}}{([\delta]_q + q^\delta)(1 - \beta e^{-i\varphi})} \quad (2.6)$$

which yields that

$$\left| \mathcal{R}_q^\delta g(z) \right| \leq \frac{([\delta]_q + q^\delta)(1 + \beta)}{[\delta]_q - ([\delta]_q + q^\delta)\beta - q^\delta e^{|z|}} \left| \mathcal{R}_q^{\delta+1}g(z) \right|. \quad (2.7)$$

Since  $\mathcal{R}_q^\delta f$  is majorized by  $\mathcal{R}_q^\delta g(z)$  in  $\mathbb{U}$ , we have

$$\mathcal{R}_q^\delta f(z) = \phi(z)\mathcal{R}_q^\delta g(z).$$

By applying  $q$ -differentiation with respect to  $z$ , we get

$$z\mathfrak{D}_q(\mathcal{R}_q^\delta f(z)) = z\mathfrak{D}_q(\phi(z))\mathcal{R}_q^\delta g(z) + z\phi(z)\mathfrak{D}_q(\mathcal{R}_q^\delta g(z)). \quad (2.8)$$

Noting the fact that Schwarz function  $\phi(z)$  satisfies the  $q$ -analogue of the result given by Nehari (cf. [7]) proved in Lemma 2.1,

$$|\mathfrak{D}_q\phi(z)| \leq \frac{1 - |\phi(z)|^2}{1 - |z|^2} \quad (2.9)$$

and using (1.14), (2.7) and (2.9) in (2.8), we have

$$|\mathcal{R}_q^{\delta+1}f(z)| \leq \left( |\phi(z)| + \left( \frac{1 - |\phi(z)|^2}{1 - |z|^2} \right) \frac{|z|q^\delta(1 + \beta)}{[\delta]_q - ([\delta]_q + q^\delta)\beta - q^\delta e^{|z|}} \right) |\mathcal{R}_q^{\delta+1}g(z)|.$$

Setting  $|z| = r$  and  $|\phi(z)| = \rho$  ( $0 \leq \rho \leq 1$ ), the above inequality leads us to the inequality

$$|\mathcal{R}_q^{\delta+1}f(z)| \leq \left( \rho + \left( \frac{1 - \rho^2}{1 - r^2} \right) \frac{rq^\delta(1 + \beta)}{[\delta]_q - ([\delta]_q + q^\delta)\beta - q^\delta e^r} \right) |\mathcal{R}_q^{\delta+1}g(z)|. \quad (2.10)$$

That is,

$$|\mathcal{R}_q^{\delta+1}f(z)| \leq \Theta_1(r, \rho)|\mathcal{R}_q^{\delta+1}g(z)|,$$

where the function  $\Theta_1(r, \rho)$  is given by

$$\Theta_1(r, \rho) = \rho + \frac{r(1 - \rho^2)q^\delta(1 + \beta)}{(1 - r^2)\{[\delta]_q - ([\delta]_q + q^\delta)\beta - q^\delta e^r\}}.$$

In order to determine the bound of  $\Theta_1(r, \rho)$ , we have to choose

$$\begin{aligned} r_1 &= \max\{r \in [0, 1) : \Theta_1(r, \rho) \leq 1, \rho \in [0, 1]\} \\ &= \max\{r \in [0, 1) : \Theta_2(r, \rho) \geq 0, \rho \in [0, 1]\}, \end{aligned}$$

where

$$\Theta_2(r, \rho) = (1 - r^2)\{[\delta]_q - ([\delta]_q + q^\delta)\beta - q^\delta e^r\} - r(1 + \rho)q^\delta(1 + \beta).$$

Obviously, for  $\rho = 1$ , the function  $\Theta_2(r, \rho)$  takes its minimum value, namely

$$\min\{\Theta_2(r, \rho) : \rho \in [0, 1]\} = \Theta_2(r, 1) = \Theta_2(r),$$

where

$$\Theta_2(r) = (1 - r^2)\{[\delta]_q - ([\delta]_q + q^\delta)\beta - q^\delta e^r\} - 2rq^\delta(1 + \beta).$$

Furthermore, if  $\Theta_2(0) = [\delta]_q > ([\delta]_q + q^\delta)\beta + q^\delta e$  and  $\Theta_2(1) = -2q^\delta(1 + \beta) < 0$ , then there exists  $r_1$  such that  $\Theta_2(r) \geq 0$  for all  $r \in [0, r_1]$ , where  $r_1 = r_1(\delta, \beta)$ , the smallest positive root of the equation (2.3). This completes the proof.  $\square$

Putting  $\beta = 0$  and  $\rho = 1$  in Theorem 2.2, we have the following corollary:

**Corollary 2.3.** *Let the function  $f \in \mathcal{A}$  and suppose that  $g \in \mathcal{RS}_q^\delta(e^z)$ . If  $\mathcal{R}_q^\delta f$  is majorized by  $\mathcal{R}_q^\delta g$  in  $\mathbb{U}$ , then*

$$|\mathcal{R}_q^{\delta+1}f(z)| \leq |\mathcal{R}_q^{\delta+1}g(z)|, \quad |z| \leq r_2, \quad (2.11)$$

where  $r_2 = r_2(\delta)$ , is the smallest positive root of the equation

$$r^2q^\delta e^r - r^2[\delta]_q - q^\delta e^r - 2rq^\delta + [\delta]_q = 0. \quad (2.12)$$

For  $\beta = 0$ ,  $q \rightarrow 1^-$  and  $\delta = 0$ , Corollary 2.3 reduces to the following result:

**Corollary 2.4.** *Let the function  $f \in \mathcal{A}$  be analytic and univalent in the open unit disk  $\mathbb{U}$  and suppose that  $g \in \mathcal{S}^*(e^z)$ . If  $f$  is majorized by  $g$  in  $\mathbb{U}$ , then*

$$|f'(z)| \leq |g'(z)| \quad (|z| \leq r_3),$$

where  $r_3$  is the smallest positive root of  $r^2e^r - 2r - e^r = 0$ .

3. A MAJORIZATION PROBLEM FOR THE CLASS  $\mathcal{R}(\mu, \tau)$ 

Due to Alitintas et al. [1], we recall the definition of the function class  $\mathcal{R}(\mu, \tau)$ , the class of functions  $h$  of the form

$$h(z) = 1 - \sum_{n=1}^{\infty} c_n z^n \quad (c_n \geq 0 ; z \in \mathbb{U}), \quad (3.1)$$

which are analytic in  $\mathbb{U}$  and satisfy the inequality

$$|h(z) + \mu z h'(z) - 1| < |\tau| \quad (\tau \in \mathbb{C} \setminus \{0\}; \Re(\mu) \geq 0).$$

Further we recall the following lemmas, which will be required in our investigation of the majorization problem for the class  $\mathcal{R}(\mu, \tau)$ .

**Lemma 3.1.** ([1]) *If the function  $h$  defined by (3.1) is in the class  $\mathcal{R}(\mu, \tau)$ , then*

$$\sum_{n=1}^{\infty} c_n \leq \frac{|\tau|}{1 + \Re(\mu)}. \quad (3.2)$$

**Lemma 3.2.** ([1]) *If the function  $h$  defined by is in the class  $\mathcal{R}(\mu, \tau)$ , then*

$$1 - \frac{|\tau|}{1 + \Re(\mu)} |z| \leq |h(z)| \leq 1 + \frac{|\tau|}{1 + \Re(\mu)} |z| \quad (z \in \mathbb{U}). \quad (3.3)$$

**Theorem 3.3.** *Let the function  $f$  and  $g$  be analytic in  $\mathbb{U}$  and suppose that the function  $g$  is normalized and also satisfies the following inclusion property:*

$$\left( \frac{z \mathcal{D}_q(\mathcal{R}_q^\delta g(z))}{\mathcal{R}_q^\delta g(z)} \right) \in \mathcal{R}(\mu, \tau).$$

*If  $\mathcal{R}_q^\delta f$  is majorized by  $\mathcal{R}_q^\delta g$  in  $\mathbb{U}$ , then*

$$|\mathcal{R}_q^{\delta+1} f(z)| \leq |\mathcal{R}_q^{\delta+1} g(z)| \quad (|z| \leq r_4), \quad (3.4)$$

*where  $r_4 = r_4(\tau, \mu, \delta)$  is the root of the cubic equation*

$$\begin{aligned} & q^\delta |\tau| r^3 - \{(q^\delta - [\delta]_q)(1 + \Re(\mu)) - 2|\tau|\} r^2 \\ & - [2(1 + \Re(\mu)) + q^\delta |\tau|] r + (q^\delta - [\delta]_q)[1 + \Re(\mu)] = 0 \end{aligned} \quad (3.5)$$

*which lies in the interval  $(0, 1)$  and  $(q^\delta - [\delta]_q)(1 + \Re(\mu)) > 0$ .*

*Proof.* For an appropriately normalized analytic function  $g$  satisfying the inclusion property, we find from the assertion of Lemma 3.2 that

$$\left| \frac{z \mathcal{D}_q(\mathcal{R}_q^\delta g(z))}{\mathcal{R}_q^\delta g(z)} \right| \geq 1 - \frac{|\tau|}{1 + \Re(\mu)} r \quad (|z| = r, 0 < r < 1) \quad (3.6)$$



or, equivalently, that

$$|\mathcal{R}_q^\delta g(z)| \leq \frac{(q^\delta + [\delta]_q)(1 + \Re(\mu) - |\tau|r)}{(q^\delta - [\delta]_q)(1 + \Re(\mu)) - q^\delta |\tau|r} |(\mathcal{R}_q^{\delta+1} g(z))| \quad (|z| = r, 0 < r < 1). \quad (3.7)$$

Since

$$\mathcal{R}_q^\delta f(z) \ll \mathcal{R}_q^\delta g(z) \quad (z \in \mathbb{U}),$$

there exists an analytic function  $\phi$  such that

$$\mathcal{R}_q^\delta f(z) = \phi(z) \mathcal{R}_q^\delta g(z) \quad \text{and} \quad |\phi(z)| < 1.$$

By applying  $q$ -differentiation with respect to  $z$ , we get

$$z \mathfrak{D}_q(\mathcal{R}_q^\delta f(z)) = z \mathfrak{D}_q(\phi(z)) \mathcal{R}_q^\delta g(z) + \phi(z) z \mathfrak{D}_q(\mathcal{R}_q^\delta g(z)). \quad (3.8)$$

Thus in view of (3.7) and using (1.14), just as in the proof of Theorem 2.2, we have

$$|\mathfrak{D}_q(\phi(z))| \leq \frac{1 - |\phi(z)|^2}{1 - |z|^2} \quad (z \in \mathbb{U})$$

and

$$|\mathfrak{D}_q(\mathcal{R}_q^\delta f(z))| \leq \left( |\phi(z)| + \frac{1 - |\phi(z)|^2}{1 - |z|^2} \cdot \frac{(1 + \Re(\mu) - |\tau|r)|z|}{(q^\delta - [\delta]_q)(1 + \Re(\mu)) - q^\delta |\tau|r} \right) |\mathfrak{D}_q(\mathcal{R}_q^\delta g(z))|,$$

$|z| = r, 0 < r < 1$ . That is,

$$\begin{aligned} |\mathcal{R}_q^{\delta+1} f(z)| &\leq \left( |\phi(z)| + \frac{1 - |\phi(z)|^2}{1 - r^2} \cdot \frac{(1 + \Re(\mu) - |\tau|r)r}{(q^\delta - [\delta]_q)(1 + \Re(\mu)) - q^\delta |\tau|r} \right) \\ &\quad \times |\mathcal{R}_q^{\delta+1} g(z)| \\ &= \Lambda_1(\rho, r) |\mathcal{R}_q^{\delta+1} g(z)|, \end{aligned} \quad (3.9)$$

where  $|z| = r, 0 < r < 1$ . We set  $|\phi(z)| = \rho$  and the function  $\Lambda_1(\rho, r)$  defined by

$$\Lambda_1(\rho, r) = \rho + \frac{1 - \rho^2}{1 - r^2} \cdot \frac{(1 + \Re(\mu) - |\tau|r)r}{(q^\delta - [\delta]_q)(1 + \Re(\mu)) - q^\delta |\tau|r}. \quad (3.10)$$

In order to determine the bound of  $\Lambda(\rho, r)$ , we have to choose

$$\begin{aligned} r_1 &= \max\{r \in [0, 1) : \Lambda_1(\rho, r) \leq 1, \rho \in [0, 1]\} \\ &= \max\{r \in [0, 1) : \Lambda_2(\rho, r) \geq 0, \rho \in [0, 1]\}, \end{aligned}$$

where, for  $0 \leq \rho \leq 1$ .

$$\Lambda_2(r, \rho) = (1 - r^2) \{ (q^\delta - [\delta]_q)(1 + \Re(\mu)) - q^\delta |\tau|r \} - r(1 + \rho)(1 + \Re(\mu) - |\tau|r).$$

Obviously, for  $\rho = 1$ , the function  $\Lambda_2(r, \rho)$  takes its minimum value, namely

$$\min\{\Lambda_2(r, \rho) : \rho \in [0, 1]\} = \Lambda_2(r, 1) = \Lambda_2(r),$$

where

$$\Lambda_2(r) = (1 - r^2)\{(q^\delta - [\delta]_q)(1 + \Re(\mu)) - q^\delta |\tau|r\} - 2r(1 + \Re(\mu) - |\tau|r).$$

Furthermore, if  $\Lambda_2(0) = (q^\delta - [\delta]_q)(1 + \Re(\mu)) > 0$  and  $\Lambda_2(1) = -2(1 + \Re(\mu) - |\tau|) < 0$ , then there exists  $r_4$  such that  $\Lambda_2(r) \geq 0$  for all  $r \in [0, r_4]$ , where  $r_4 = r_4(\tau, \mu, \delta)$ , the smallest positive root of the equation (3.5) which completes the proof of Theorem 3.3.  $\square$

**Remark 3.4.** Specializing the parameters  $\delta, \beta$  in (1.15) one can define the various other interesting subclasses of  $\mathcal{RS}_q^\delta(\beta, e^z)$ , involving  $q$ -calculus operator and one can easily derive the result as in Theorem 2.2. Further as mentioned in [11] we can define new subclasses  $\mathcal{RS}_q^\delta(\beta, 1 + \sin z)$ ,  $\mathcal{RS}_q^\delta(\beta, \cos z)$ , and  $\mathcal{RS}_q^\delta(\beta, z + \sqrt{1 + z^2})$ , and investigate a majorization problem for these classes.

**Acknowledgments:** This work was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology(2019R111A3A01050861).

#### REFERENCES

- [1] O. Altıntaş, Ö. Özkan and H.M. Srivastava, *Majorization by starlike functions of complex order*, Complex Variables Theory Appl., **46** (2001), 207–218.
- [2] S.P. Goyal and P. Goswami, *Majorization for certain classes of analytic functions defined by fractional derivatives*, Appl. Math. Lett., **22** (2009), 1855–1858.
- [3] F.H. Jackson, *On  $q$ -functions and a certain difference operator*, Trans. Royal Soc. Edinburgh, **46** (1908), 253–281.
- [4] S. Kanas and D. Răducanu, *Some subclass of analytic functions related to conic domains*, Math. Slovaca, **64**(5) (2014), 1183–1196.
- [5] T.H. MacGregor, *Majorization by univalent functions*, Duke Math. J., **34** (1967), 95–102.
- [6] S.S. Miller and P.T. Mocanu, *Differential subordinations* Monographs and Textbooks in Pure and Applied Mathematics, 225, Dekker, New York, 2000.
- [7] Z. Nehari, *Conformal mapping* MacGra-Hill Book Company, New York; Toronto and London, 1952.
- [8] R. Mendiratta, S. Nagpal and V. Ravichandran, *On a subclass of strongly starlike functions associated with exponential function*, Bull. Malaysian Math. Soc., **38** (2014), 365–386.
- [9] S. Ruscheweyh, *New criteria for univalent functions*, roc. Amer. Math. Soc., **49** (1975), 109–115.
- [10] K.A. Selvakumaran, S.D. Purohit and A. Secer, *Majorization for a class of analytic functions defined by  $q$ -differentiation*, Math. Prob. Engineering, **2014**, Article ID: 653917, 5 pages.
- [11] H. Tang, H.M. Srivastava, S.H. Li and G.T. Deng, *Majorization results for subclasses of starlike functions based on the sine and cosine functions*, Bull. Iran. Math. Soc., **46** (2020), 389–391.

- [12] H. Tang and G. Deng, *Majorization problems for certain classes of multivalent analytic functions related with the Srivastava-khairnar-more operator and exponential function*, *Filomat*, **32**(15) (2018), 5319–5328.