



CONVERGENCE THEOREM FOR A GENERALIZED φ -WEAKLY CONTRACTIVE NONSELF MAPPING IN METRICALLY CONVEX METRIC SPACES

Kyung Soo Kim

Department of Mathematics Education, Kyungnam University
Changwon, Gyeongnam, 51767, Republic of Korea
e-mail: kksmj@kyungnam.ac.kr

Abstract. A convergence theorem for a generalized φ -weakly contractive mapping is proved which satisfy a generalized contraction condition on a complete metrically convex metric space. The result in this paper generalizes the relevant results due to Rhoades [18], Alber and Guerre-Delabriere [1], Khan and Imdad [14], Xue [20] and others. An illustrative example is also furnished in support of our main result.

1. INTRODUCTION AND PRELIMINARIES

Let (X, d) be a metric space. A mapping $T : X \rightarrow X$ is said to be a *contraction* if there exists a constant $\alpha \in (0, 1)$ such that

$$d(Tx, Ty) \leq \alpha \cdot d(x, y), \quad \forall x, y \in X.$$

Theorem 1.1. ([4]) *Let (X, d) be a complete metric space and let $T : X \rightarrow X$ be a contraction mapping. Then T has a unique fixed point.*

There are so many extensions and generalizations of the theorem mentioned above (see [5]-[9]).

A mapping $T : X \rightarrow X$ is said to be a *φ -weak contraction* if there exists a continuous and nondecreasing function $\varphi : [0, \infty) \rightarrow [0, \infty)$ with $\varphi^{-1}(0) = \{0\}$

⁰Received December 30, 2020. Revised March 11, 2021. Accepted March 15, 2021.

⁰2010 Mathematics Subject Classification: 49J40, 47H10, 47H17.

⁰Keywords: Metrically convex metric space, generalized φ -weakly contractive nonself mapping, fixed point.

and $\lim_{t \rightarrow \infty} \varphi(t) = \infty$ such that

$$d(Tx, Ty) \leq d(x, y) - \varphi(d(x, y)), \quad \forall x, y \in X. \quad (1.1)$$

If X is bounded, then the infinity condition can be omitted.

The concept of the φ -weak contraction was introduced by Alber and Guerre-Delabriere [1] in 1997, who proved the existence of fixed points in Hilbert spaces. Later Rhoades [18] in 2001 extended the results of [1] to metric spaces.

Theorem 1.2. ([18]) *Let (X, d) be a complete metric space and $T : X \rightarrow X$ be a φ -weakly contractive self-map on X . Then T has a unique fixed point p in X .*

Remark 1.3. Theorem 1.2 is one of generalizations of the Banach contraction principle because it takes $\varphi(t) = (1 - \alpha)t$ for $\alpha \in (0, 1)$, then φ -weak contraction contains contraction as special cases.

In 2016, Xue [20] introduced a new contraction type mapping as follows.

Definition 1.4. ([20]) A mapping $T : X \rightarrow X$ is said to be a *generalized φ -weak contraction* if there exists a continuous and nondecreasing function $\varphi : [0, \infty) \rightarrow [0, \infty)$ with $\varphi(0) = 0$ such that

$$d(Tx, Ty) \leq d(x, y) - \varphi(d(Tx, Ty)), \quad \forall x, y \in X \quad (1.2)$$

holds.

We notice immediately that if $T : X \rightarrow X$ is a φ -weak contraction, then T is a generalized φ -weak contraction. However, the converse is not true in general.

Example 1.5. ([15]) Let $X = (-\infty, +\infty)$ be endowed with the Euclidean metric $d(x, y) = |x - y|$ and let $Tx = \frac{2}{5}x$ for each $x \in X$.

Define $\varphi(t) : [0, +\infty) \rightarrow [0, +\infty)$ by $\varphi(t) = \frac{4}{3}t$. Then T satisfies (1.2), but T does not satisfy inequality (1.1). Indeed,

$$\begin{aligned} d(Tx, Ty) &= \left| \frac{2}{5}x - \frac{2}{5}y \right| \\ &\leq |x - y| - \frac{4}{3} \cdot \frac{2}{5} |x - y| \\ &= d(x, y) - \varphi(d(Tx, Ty)) \end{aligned}$$

and

$$\begin{aligned} d(Tx, Ty) &= \left| \frac{2}{5}x - \frac{2}{5}y \right| \\ &\geq |x - y| - \frac{4}{3}|x - y| \\ &= d(x, y) - \varphi(d(x, y)) \end{aligned}$$

for all $x, y \in X$.

Example 1.6. ([20]) Let $X = [0, +\infty)$ be endowed by $d(x, y) = |x - y|$ and let $Tx = \frac{x}{1+x}$ for each $x \in X$. Define $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ by $\varphi(t) = \frac{t^2}{1+t}$. Then

$$\begin{aligned} d(Tx, Ty) &= \left| \frac{x}{1+x} - \frac{y}{1+y} \right| = \frac{|x - y|}{(1+x)(1+y)} \\ &\leq \frac{|x - y|}{1 + |x - y|} = |x - y| - \frac{|x - y|^2}{1 + |x - y|} \\ &= d(x, y) - \varphi(d(x, y)) \end{aligned}$$

holds for all $x, y \in X$. So T is a φ -weak contraction. However, T is not a contraction.

Remark 1.7. The above examples show that the class of generalized φ -weak contractions properly includes the class of φ -weak contractions and the class of φ -weak contractions properly includes the class of contractions. In fact, let $T : X \rightarrow X$ be a contraction, there exists $\alpha \in (0, 1)$ such that

$$d(Tx, Ty) \leq \alpha \cdot d(x, y), \quad \forall x, y \in X.$$

Then

$$\begin{aligned} d(Tx, Ty) &\leq \alpha \cdot d(x, y) = d(x, y) - (1 - \alpha)d(x, y) \\ &= d(x, y) - \varphi(d(x, y)), \end{aligned}$$

where, $\varphi(d(x, y)) = (1 - \alpha)d(x, y)$. So, T is a φ -weak contraction. Moreover, let T be a φ -weak contraction, from property of φ , we have $d(Tx, Ty) \leq d(x, y)$ and

$$\varphi(d(Tx, Ty)) \leq \varphi(d(x, y)).$$

From (1.1),

$$\begin{aligned} d(Tx, Ty) &\leq d(x, y) - \varphi(d(x, y)) \\ &\leq d(x, y) - \varphi(d(Tx, Ty)), \quad \forall x, y \in X. \end{aligned}$$

Therefore, T is a generalized φ -weak contraction.

In many applications the mappings involved is not always a self map. So, it is interested to find sufficient conditions for such mappings will guarantee the existence of a fixed point. Assad and Kirk [3] initiated the study of fixed point of nonself mappings in metrically convex spaces. The technique due to Assad and Kirk [3] has been utilized by many researchers and there exists considerable literature on this topic. To mentioned a few, we cite [2], [10]-[17], [19], [21].

In this paper, we prove a convergence theorem for single valued nonself mapping by utilizing the idea of Rhoades [18] which either partially or completely generalize the results due to Rhoades [18], Alber and Guerre-Delabriere [1], Khan and Imdad [14] and others.

Definition 1.8. Let K be a nonempty subset of a metric space (X, d) . A mapping $T : K \rightarrow X$ is said to be a generalized φ -weak contraction if $Tx \in K$, $Tx \cap K$ is nonempty and

$$d(Tx, Ty) \leq d(x, y) - \varphi(d(Tx, Ty)), \quad (1.3)$$

where $\varphi : [0, \infty) \rightarrow [0, \infty)$ is continuous and nondecreasing function with $\varphi(t) = 0$ iff $t = 0$.

Definition 1.9. ([3]) A metric space (X, d) is said to be metrically convex if for any $x, y \in X$ with $x \neq y$ there exists a point $z \in X$, $x \neq z \neq y$ such that

$$d(x, z) + d(z, y) = d(x, y).$$

Remark 1.10. ([3]) If K is a closed subset of the complete and convex metric space X and if $x \in K$, $y \notin K$, then there exists a point z in the boundary of K such that

$$d(x, z) + d(z, y) = d(x, y).$$

We use the symbol ∂K to denote the boundary of K .

2. MAIN RESULT

In this section, we will prove convergence theorem for a single valued generalized φ -weak contractive nonself mapping.

Theorem 2.1. *Let (X, d) be a complete metrically convex metric space and K be a nonempty closed subset of X . Let $T : K \rightarrow X$ be a generalized φ -weak contraction defined as (1.3) for each $x \in \partial K$ and $Tx \in K$. Then T has a unique fixed point in K .*

Proof. We proceed to construct two sequences $\{x_n\}$ and $\{y_n\}$ in the following way. Let $x_0 \in K$. Define $y_1 = Tx_0$. If $y_1 \in K$, set $y_1 = x_1$. If $y_1 \notin K$, then choose $x_1 \in \partial K$ such that

$$d(x_0, x_1) + d(x_1, y_1) = d(x_0, y_1).$$

Define $y_2 = Tx_1$. If $y_2 \in K$, set $y_2 = x_2$. If $y_2 \notin K$, then choose $x_2 \in \partial K$ such that

$$d(x_1, x_2) + d(x_2, y_2) = d(x_1, y_2).$$

Continuing this process, we obtain two sequences $\{x_n\}$ and $\{y_n\}$ such that

$$\begin{aligned} & \text{(i) } y_{n+1} = Tx_n, \\ & \text{(ii) } \begin{cases} x_n = y_n, & \text{if } y_n \in K, \\ \text{choose } x_n \in \partial K \text{ such that} \\ \quad d(x_{n-1}, x_n) + d(x_n, y_n) = d(x_{n-1}, y_n), & \text{if } y_n \notin K. \end{cases} \end{aligned}$$

Here, we have two types of sets we denote as follows:

$$\begin{aligned} P &= \{x_i \in \{x_n\} : x_i = y_i, i = 1, 2, \dots\} \\ &= \{x_i \in \{x_n\} : y_i = Tx_{i-1} \in K, i = 1, 2, \dots\} \end{aligned}$$

and

$$\begin{aligned} Q &= \{x_i \in \{x_n\} : x_i \neq y_i, i = 1, 2, \dots\} \\ &= \{x_i \in \{x_n\} : y_i = Tx_{i-1} \notin K, i = 1, 2, \dots\} \\ &= \{x_i \in \{x_n\} : d(x_{i-1}, x_i) + d(x_i, y_i) = d(x_{i-1}, y_i), x_i \in \partial K, i = 1, 2, \dots\}. \end{aligned}$$

Observe that if $x_n \in Q$ for some n , then x_{n-1} and $x_{n+1} \in P$.

Now, we wish to estimate $d(x_n, x_{n+1})$. Now we distinguish the following cases.

Case 1. If x_n and $x_{n+1} \in P$, then

$$\begin{aligned} d(x_n, x_{n+1}) &= d(y_n, y_{n+1}) = d(Tx_{n-1}, Tx_n) \\ &\leq d(x_{n-1}, x_n) - \varphi(d(x_n, x_{n+1})) \\ &\leq d(x_{n-1}, x_n), \end{aligned}$$

by monotonicity of φ .

Case 2. If $x_n \in P$ and $x_{n+1} \in Q$, then

$$d(x_n, x_{n+1}) + d(x_{n+1}, y_{n+1}) = d(x_n, y_{n+1}).$$

Therefore

$$\begin{aligned} d(x_n, x_{n+1}) &\leq d(x_n, y_{n+1}) = d(y_n, y_{n+1}) = d(Tx_{n-1}, Tx_n) \\ &\leq d(x_{n-1}, x_n) - \varphi(d(y_n, y_{n+1})) \\ &\leq d(x_{n-1}, x_n). \end{aligned}$$

Case 3. If $x_n \in Q$ and $x_{n+1} \in P$. Since $x_n \in Q$ and $x_{n-1} \in P$, we have $x_{n-1} = y_{n-1}$. It follows

$$\begin{aligned} d(x_n, x_{n+1}) &\leq d(x_n, y_n) + d(y_n, x_{n+1}) = d(x_n, y_n) + d(y_n, y_{n+1}) \\ &= d(x_n, y_n) + d(Tx_{n-1}, Tx_n) \\ &\leq d(x_n, y_n) + d(x_{n-1}, x_n) - \varphi(d(y_n, y_{n+1})) \\ &\leq d(x_{n-1}, y_n) = d(y_{n-1}, y_n) = d(Tx_{n-2}, Tx_{n-1}) \\ &\leq d(x_{n-2}, x_{n-1}) - \varphi(d(y_{n-1}, y_n)) \\ &\leq d(x_{n-2}, x_{n-1}). \end{aligned}$$

Case 4. The only other possibility, $x_n \in Q$ and $x_{n+1} \in Q$, cannot occur.

Thus in all the cases, we have

$$\begin{aligned} d(x_n, x_{n+1}) &\leq \max\{d(x_{n-1}, x_n) - \varphi(d(y_n, y_{n+1})), d(x_{n-2}, x_{n-1}) - \varphi(d(y_{n-1}, y_n))\} \\ &\leq \max\{d(x_{n-1}, x_n), d(x_{n-2}, x_{n-1})\}. \end{aligned} \quad (2.1)$$

By (2.1), it follows that the sequence $\{d(x_n, x_{n+1})\}$ is monotone decreasing. Hence

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0. \quad (2.2)$$

Next, we prove that the sequence $\{x_n\}$ is Cauchy. Suppose that the sequence $\{x_n\}$ is not Cauchy. Then there exists $\varepsilon > 0$ for which we can find subsequences $\{x_{n_k}\}$ and $\{x_{n_l}\}$ such that

$$d(x_{n_k}, x_{n_l}) \geq \varepsilon.$$

Corresponding to each n_k , we can find n_l in such a way that the smallest positive integer $n_l > n_k$ satisfying

$$d(x_{n_k}, x_{n_l-1}) < \varepsilon.$$

Then we have

$$\varepsilon \leq d(x_{n_k}, x_{n_l}) \leq d(x_{n_k}, x_{n_l-1}) + d(x_{n_l-1}, x_{n_l}) < \varepsilon + d(x_{n_l-1}, x_{n_l}).$$

Letting $n \rightarrow \infty$, we have

$$\lim_{n \rightarrow \infty} d(x_{n_k}, x_{n_l}) = \varepsilon. \quad (2.3)$$

By (2.1), if $d(x_{n-1}, x_n) - \varphi(d(y_n, y_{n+1})) \geq d(x_{n-2}, x_{n-1}) - \varphi(d(y_{n-1}, y_n))$, then

$$d(x_n, x_{n+1}) \leq d(x_{n-1}, x_n) - \varphi(d(y_n, y_{n+1})).$$

So we have

$$\begin{aligned} d(x_{n_k}, x_{n_l}) &\leq d(x_{n_k}, x_{n_{k+1}}) + d(x_{n_{k+1}}, x_{n_{k+2}}) + \cdots + d(x_{n_{l-1}}, x_{n_l}) \\ &\leq d(x_{n_{k-1}}, x_{n_k}) - \varphi(d(y_{n_k}, y_{n_{k+1}})) \\ &\quad + d(x_{n_k}, x_{n_{k+1}}) - \varphi(d(y_{n_{k+1}}, y_{n_{k+2}})) \\ &\quad \vdots \\ &\quad + d(x_{n_{l-2}}, x_{n_{l-1}}) - \varphi(d(y_{n_{l-1}}, y_{n_l})). \end{aligned} \tag{2.4}$$

If $d(x_{n-1}, x_n) - \varphi(d(y_n, y_{n+1})) \leq d(x_{n-2}, x_{n-1}) - \varphi(d(y_{n-1}, y_n))$, then

$$d(x_n, x_{n+1}) \leq d(x_{n-2}, x_{n-1}) - \varphi(d(y_{n-1}, y_n)).$$

So we get

$$\begin{aligned} d(x_{n_k}, x_{n_l}) &\leq d(x_{n_k}, x_{n_{k+1}}) + d(x_{n_{k+1}}, x_{n_{k+2}}) + \cdots + d(x_{n_{l-1}}, x_{n_l}) \\ &\leq d(x_{n_{k-2}}, x_{n_{k-1}}) - \varphi(d(y_{n_{k-1}}, y_{n_k})) \\ &\quad + d(x_{n_{k-1}}, x_{n_k}) - \varphi(d(y_{n_k}, y_{n_{k+1}})) \\ &\quad \vdots \\ &\quad + d(x_{n_{l-3}}, x_{n_{l-2}}) - \varphi(d(y_{n_{l-2}}, y_{n_{l-1}})). \end{aligned} \tag{2.5}$$

From (2.2), (2.3) and $\varepsilon > 0$, on letting $n \rightarrow \infty$ in (2.4) and (2.5), we have a contradiction. Thus the sequence $\{x_n\}$ is Cauchy and since X is complete and K is closed, $\{x_n\}$ converges to a point $p \in K$. Also observe that there exists a subsequence $\{x_{n_j}\}$ of $\{x_n\}$ each of whose terms is in the set P , that is

$$x_{n_j} = y_{n_j}, \quad j = 1, 2, \dots.$$

Thus by condition (i),

$$y_{n_j} = Tx_{n_{j-1}}, \quad j = 1, 2, \dots.$$

Since $x_{n_{j-1}} \rightarrow p$ as $j \rightarrow \infty$, we have $Tx_{n_{j-1}} \rightarrow Tp$ as $j \rightarrow \infty$, that is,

$$\lim_{j \rightarrow \infty} y_{n_j} = Tp.$$

Now, we will show that p is a fixed point of T . Suppose that $p \neq Tp$. Then $d(Tp, p) > 0$. So we get

$$\begin{aligned} d(Tp, x_{n_j+1}) &= d(Tp, y_{n_j+1}) = d(Tp, Tx_{n_j}) \\ &\leq d(p, x_{n_j}) - \varphi(d(Tp, x_{n_j+1})), \end{aligned}$$

that is,

$$d(Tp, x_{n_j+1}) + \varphi(d(Tp, x_{n_j+1})) \leq d(p, x_{n_j}).$$

Taking the limit as $j \rightarrow \infty$ to above inequality,

$$d(Tp, p) + \varphi(d(Tp, p)) \leq d(p, p) = 0.$$

Since $d(Tp, p) > 0$, this is a contradiction. Therefore

$$Tp = p.$$

Finally, we will prove that the uniqueness of fixed point. Assume that p and q are two fixed points of T . Then

$$\begin{aligned} d(p, q) &= d(Tp, Tq) \\ &\leq d(p, q) - \varphi(d(Tp, Tq)) \\ &= d(p, q) - \varphi(d(p, q)), \end{aligned}$$

that is,

$$\varphi(d(p, q)) \leq 0.$$

By the condition of φ , we have

$$d(p, q) = 0.$$

Therefore

$$p = q.$$

This completes the proof. \square

Remark 2.2. By setting $K = X$ in Theorem 2.1, one deduces a theorem due to Xue [20].

3. AN ILLUSTRATIVE EXAMPLE

In this section, we furnish an example to establish the utility of our result.

Example 3.1. Let $X = \mathbb{R}$ with Euclidean metric and $K = [0, 1]$. Define $T : K \rightarrow X$ and $\varphi : [0, \infty) \rightarrow [0, \infty)$ by

$$Tx = \frac{2}{5}(x - x^2), \quad \varphi(t) = \frac{4}{3}t.$$

Since boundary of K , $\partial K = \{0, 1\}$, we have

$$T0 = 0 \in K, \quad T1 = 0 \in K.$$

For the verification of generalized φ -weak contraction condition (1.3) the following cases arise:

$$\begin{aligned}
d(Tx, Ty) &= |Tx - Ty| \\
&= \frac{2}{5}|(x - y) - (x - y)(x + y)| \\
&= \frac{2}{5}|(x - y)| \cdot |1 - x - y| \\
&\leq |x - y| - |x - y| \cdot |1 - x - y| + \frac{2}{5}|(x - y)| \cdot |1 - x - y| \\
&= |x - y| - \frac{3}{5}|x - y| \cdot |1 - x - y| \\
&< |x - y| - \frac{8}{15}|x - y| \cdot |1 - x - y| \\
&= |x - y| - \frac{4}{3} \cdot \frac{2}{5}|x - y| \cdot |1 - x - y| \\
&= d(x, y) - \varphi(d(Tx, Ty)) \quad \forall x, y \in K.
\end{aligned}$$

Thus the generalized φ -weak contraction condition (1.3) and all other conditions of the Theorem 2.1 are satisfied. Moreover, 0 is the unique fixed point of T .

Acknowledgments: The author would like to thank the referees for their valuable comments and suggestions which improved the presentation of this paper. This work was supported by Kyungnam University Foundation Grant, 2020.

REFERENCES

- [1] Ya.I. Alber and S. Guerre-Delabriere, *Principle of weakly contractive maps in Hilbert spaces*, in new results in Operator theory, Advances and Applications (Ed. by I. Gohberg and Y. Lyubich), Birkhauser Verlag Basel, **98** (1997), 7–22.
- [2] N.A. Assad, *On a fixed point theorem of Kannan in Banach spaces*, Tamkang J. Math., **7** (1976), 91–94.
- [3] N.A. Assad and W.A. Kirk, *Fixed point theorems for set-valued mappings of contractive type*, Pacific J. Math., **43**(3) (1972), 553–562.
- [4] S. Banach, *Sur les operations dans les ensembles abstraits et leur application aux equations integrales*, Fundam. Math., **3** (1922), 133–181.
- [5] I. Beg and M. Abbas, *Coincidence point and invariant approximation for mappings satisfying generalized weak contractive condition*, Fixed Point Theory and Appl., Article ID (74503), **2006** (2006), 1–7.
- [6] D.W. Boyd and J.S.W. Wong, *On nonlinear contractions*, Proc. Amer. Math. Soc., **20** (1969), 458–464.
- [7] C.E. Chidume, H. Zegeye and S.J. Aneke, *Approximation of fixed points of weak contractive nonself maps in Banach spaces*, J. Math. Anal. Appl., **270**(1) (2002), 189–199.
- [8] P.N. Dutta and B.S. Choudhury, *A generalisation of contraction principle in metric spaces*, Fixed Point Theory and Appl., Article ID (406368) **2008** (2008), 1–8.
- [9] M. Edelstein, *On fixed and periodic points under contractive mappings*, J. London Math. Soc., **37** (1962), 74–79.

- [10] M. Imdad, *Remarks on fixed point theorems for nonself mappings*, Aligarh Bull. Math., **22**(1) (2003), 15–17.
- [11] M. Imdad and L. Khan, *Common fixed point theorems for a pair of non-self mappings*, Nonlinear Anal. Forum, **10**(1) (2005), 21–35.
- [12] M. Imdad and L. Khan, *Common fixed point theorems for two pairs of non-self mappings*, J. Appl. Math. Computing, **21**(1-2) (2006), 269–287.
- [13] L. Khan, *Fixed point theorem for weakly contractive maps in metrically convex spaces under C -class function*, Nonlinear Funct. Anal. Appl., **25**(1) (2020), 153–160.
- [14] L. Khan and M. Imdad, *Fixed point theorem for weakly contractive maps in metrically convex spaces*, Nonlinear Funct. Anal. Appl., **21**(4) (2016), 685–691.
- [15] K.S. Kim, *Equivalence between some iterative schemes for generalized φ -weak contraction mappings in $CAT(0)$ spaces*, East Asian Math. J., **33**(1) (2017), 11–22.
- [16] K.S. Kim, *Convergence and stability of generalized φ -weak contraction mapping in $CAT(0)$ spaces*, Open Math., **15** (2017), 1063–1074, <https://doi.org/10.1515/math-2017-0089>.
- [17] K.S. Kim, *Best proximity point of contraction type mapping in metric space*, J. Comput. Anal. Appl., **27**(6) (2019), 1023–1033.
- [18] B.E. Rhoades, *Some theorems on weakly contractive maps*, Nonlinear Anal., **47**(4) (2001), 2683–2693.
- [19] B.E. Rhoades, *A fixed point theorem for some nonself mappings*, Math. Japonica, **23**(4) (1978), 457–459.
- [20] Z. Xue, *The convergence of fixed point for a kind of weak contraction*, Nonlinear Funct. Anal. Appl., **21**(3) (2016), 497–500.
- [21] Q. Zhang and Y. Song, *Fixed point theory for generalized φ -weak contractions*, Appl. Math. Lett., **22** (2009), 75–78.