



NUMERICAL SOLUTIONS OF NONLINEAR VOLTERRA-FREDHOLM INTEGRO-DIFFERENTIAL EQUATIONS BY USING MADM AND VIM

Ayoob M. Abed¹, Muhammed F. Younis² and Ahmed A. Hamoud³

¹Department of Mathematics, Thi Qar Directorates of Education
Ministry of Education, Thi Qar, Iraq
e-mail: ayob.moha75@gmail.com

²Department of Mathematics, Thi Qar Directorates of Education
Ministry of Education, Thi Qar, Iraq
e-mail: muhammedfaleh@yahoo.com

³Department of Mathematics, Faculty of Education and Science
Taiz University, Taiz-380 015, Yemen
e-mail: ahmed.hamoud@taiz.edu.ye

Abstract. The aim of the current work is to investigate the numerical study of a nonlinear Volterra-Fredholm integro-differential equation with initial conditions. Our approximation techniques modified adomian decomposition method (MADM) and variational iteration method (VIM) are based on the product integration methods in conjunction with iterative schemes. The convergence of the proposed methods have been proved. We conclude the paper with numerical examples to illustrate the effectiveness of our methods.

1. INTRODUCTION

Integral and integro-differential equations are important in many applications of applied mathematics. Since many physical problems are modeled by integro-differential equations, the numerical solutions of such integro-differential equations have been highly studied by many authors. In recent

⁰Received August 21, 2021. Revised November 4, 2021. Accepted December 10, 2021.

⁰2020 Mathematics Subject Classification: 49M27, 65K10, 45J05, 65R20.

⁰Keywords: Nonlinear Volterra-Fredholm integro-differential equation, modified adomian decomposition method, variational iteration method, convergence theorem.

⁰Corresponding author: Ahmed A. Hamoud (ahmed.hamoud@taiz.edu.ye).

years, numerous works have been focusing on the development of more advanced and efficient methods for integral equations and integro-differential equations, for example [2, 9, 13, 16, 19, 20, 21, 23].

In recent years, many authors focus on the development of numerical and analytical techniques for integro-differential equations. For instance, we can remember the following works. Abbasbandy and Elyas [1] studied some applications on VIM for solving system of nonlinear Volterra integro-differential equations, Alao et al. [3] used ADM and VIM for solving integro-differential equations, Salih and Mehmet [28] applied the Taylor method to solve the linear Volterra-Fredholm integro-differential equations, Mittal and Nigam [25] applied the ADM to approximate solutions for fractional integro-differential equations, and Behzadi et al. [5] solved some class of nonlinear Volterra-Fredholm integro-differential equations by HAM. Moreover, several authors have applied the ADM and VIM to find the approximate solutions of various types of integro-differential equations [4, 6, 7, 8, 14, 15, 25].

In this paper, we consider nonlinear Volterra-Fredholm integro-differential equation of the form:

$$\begin{aligned} \sum_{j=0}^k \xi_j(x) u^{(j)}(x) &= f(x) + \lambda_1 \int_a^x K_1(x, t) G_1(u(t)) dt \\ &+ \lambda_2 \int_a^b K_2(x, t) G_2(u(t)) dt, \end{aligned} \quad (1.1)$$

with the initial conditions

$$u^{(r)}(a) = b_r, \quad r = 0, 1, 2, \dots, (k-1), \quad (1.2)$$

where $u^{(j)}(x)$ is the j^{th} derivative of the unknown function $u(x)$ that will be determined, $K_i(x, t)$, $i = 1, 2$ are the kernels of the equation, $f(x)$ and $\xi_j(x)$ are analytic functions, G_1 and G_2 are nonlinear functions of u and $a, b, \lambda_1, \lambda_2$, and b_r are real finite constants.

To obtain the approximate solution, we integrating (k)-times in the interval $[a, x]$ with respect to x we obtain,

$$\begin{aligned} u(x) &= L^{-1} \left(\frac{f(x)}{\xi_k(x)} \right) + \sum_{r=0}^{k-1} \frac{1}{r!} (x-a)^r b_r + \lambda_1 L^{-1} \left(\int_a^x \frac{K_1(x, t)}{\xi_k(x)} G_1(u(t)) dt \right) \\ &+ \lambda_2 L^{-1} \left(\int_a^b \frac{K_2(x, t)}{\xi_k(x)} G_2(u(t)) dt \right) - \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} u_n^{(j)}(x) \right), \end{aligned} \quad (1.3)$$

where L^{-1} is the multiple integration operator given as follows:

$$L^{-1}(\cdot) = \int_a^x \int_a^x \cdots \int_a^x (\cdot) dt dt \cdots dt \quad (k - \text{times}).$$

2. ANALYSIS OF THE METHODS

Some powerful methods have been focusing on the development of more advanced and efficient methods for solving integro-differential equations such as the MADM [2, 4, 6, 8, 14, 18, 22, 24, 25, 26, 29] and VIM [3, 7, 11, 12, 10].

2.1. Modified Adomian Decomposition Method (MADM). The modified decomposition method will facilitate the computational process and further accelerate the convergence of the series solution. The MADM depends mainly on splitting the function $f(x)$ into two parts.

$$f(x) = f_1(x) + f_2(x). \tag{2.1}$$

From Eq. (1.3) and Eq. (2.1), we obtain

$$\begin{aligned} u(x) = & L^{-1} \left(\frac{f_1(x)}{\xi_k(x)} \right) + L^{-1} \left(\frac{f_2(x)}{\xi_k(x)} \right) + \sum_{r=0}^{k-1} \frac{1}{r!} (x-a)^r b_r \\ & + \lambda_1 L^{-1} \left(\int_a^x \frac{K_1(x,t)}{\xi_k(x)} G_1(u(t)) dt \right) \\ & + \lambda_2 L^{-1} \left(\int_a^b \frac{K_2(x,t)}{\xi_k(x)} G_1(u(t)) dt \right) - \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} u_n^{(j)}(x) \right). \end{aligned}$$

We assume

$$G_1(u(x)) = \sum_{n=0}^{\infty} A_n, \quad G_2(u(x)) = \sum_{n=0}^{\infty} B_n, \tag{2.2}$$

where $A_n, B_n, n \geq 0$ are the adomian polynomials determined formally as follows:

$$A_n = \frac{1}{n!} \left[\frac{d^n}{d\mu^n} G_1 \left(\sum_{i=0}^{\infty} \mu^i u_i \right) \right]_{\mu=0}, \quad B_n = \frac{1}{n!} \left[\frac{d^n}{d\mu^n} G_2 \left(\sum_{i=0}^{\infty} \mu^i u_i \right) \right]_{\mu=0}. \tag{2.3}$$

The adomian polynomials were introduced in [17, 18, 29] as:

$$\begin{aligned} A_0 &= G_1(u_0); \\ A_1 &= u_1 G_1'(u_0); \\ A_2 &= u_2 G_1'(u_0) + \frac{1}{2!} u_1^2 G_1''(u_0); \\ A_3 &= u_3 G_1'(u_0) + u_1 u_2 G_1''(u_0) + \frac{1}{3!} u_1^3 G_1'''(u_0), \dots \end{aligned}$$

and

$$\begin{aligned} B_0 &= G_2(u_0); \\ B_1 &= u_1 G_2'(u_0); \\ B_2 &= u_2 G_2'(u_0) + \frac{1}{2!} u_1^2 G_2''(u_0); \\ B_3 &= u_3 G_2'(u_0) + u_1 u_2 G_2''(u_0) + \frac{1}{3!} u_1^3 G_2'''(u_0), \dots \end{aligned}$$

The components u_0, u_1, u_2, \dots are usually determined recursively by

$$\begin{aligned} u_0 &= L^{-1} \left(\frac{f_1(x)}{\xi_k(x)} \right) + \sum_{r=0}^{k-1} \frac{1}{r!} (x-a)^r b_r, \\ u_1 &= L^{-1} \left(\frac{f_2(x)}{\xi_k(x)} \right) + \lambda_1 L^{-1} \left(\int_a^x \frac{K_1(x,t)}{\xi_k(x)} A_0(t) dt \right) \\ &\quad + \lambda_2 L^{-1} \left(\int_a^b \frac{K_2(x,t)}{\xi_k(x)} B_0(t) dt \right) - \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} u_0^{(j)}(x) \right), \\ u_{n+1} &= \lambda_1 L^{-1} \left(\int_a^x \frac{K_1(x,t)}{\xi_k(x)} A_n(t) dt \right) + \lambda_2 L^{-1} \left(\int_a^b \frac{K_2(x,t)}{\xi_k(x)} B_n(t) dt \right) \\ &\quad - \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} u_n^{(j)}(x) \right), \quad n \geq 1. \end{aligned}$$

Then, $u(x) = \sum_{i=0}^n u_i(x)$ as the approximate solution.

2.2. Variational Iteration Method (VIM). We consider the following general differential equation:

$$Lu(t) + Nu(t) = f(t),$$

where L is a linear operator, N is a nonlinear operator and $f(t)$ is inhomogeneous term. According to variational iteration method [3, 10, 27], the terms of a sequence $\{u_n\}$ are constructed such that this sequence converges to the exact solution. The terms u_n are calculated by a correction functional as follows:

$$u_{n+1}(t) = u_n(t) + \int_0^t \lambda(\tau) (Lu_n(\tau) + N\tilde{u}(\tau) - f(\tau)) d\tau. \quad (2.4)$$

To obtain the approximation solution of IVP (1.1) – (1.2), according to the VIM, the iteration formula (2.4) can be written as follows:

$$u_{n+1}(x) = u_n(x) + L^{-1} \left[\lambda(x) \left[\sum_{j=0}^k \xi_j(x) u_n^{(j)}(x) - f(x) - \lambda_1 \int_a^x K_1(x,t) G_1(u_n(t)) dt - \lambda_2 \int_a^b K_2(x,t) G_2(u_n(t)) dt \right] \right].$$

To find the optimal $\lambda(x)$, we proceed as follows:

$$\begin{aligned} \delta u_{n+1}(x) &= \delta u_n(x) + \delta L^{-1} \left[\lambda(x) \left[\sum_{j=0}^k \xi_j(x) u_n^{(j)}(x) - f(x) - \lambda_1 \int_a^x K_1(x,t) G_1(u_n(t)) dt - \lambda_2 \int_a^b K_2(x,t) G_2(u_n(t)) dt \right] \right] \\ &= \delta u_n(x) + \lambda(x) \delta u_n(x) - L^{-1} \left[\delta u_n(x) \lambda'(x) \right]. \end{aligned} \tag{2.5}$$

From Eq.(2.5), the stationary conditions can be obtained as follows:

$$\lambda'(x) = 0, \text{ and } 1 + \lambda(x)|_{x=t} = 0.$$

As a result, the Lagrange multipliers can be identified as $\lambda(x) = -1$, and by substituting in Eq.(2.5), the following iteration formula is obtained:

$$\begin{aligned} u_0(x) &= L^{-1} \left[\frac{f(x)}{\xi_k(x)} \right] + \sum_{r=0}^{k-1} \frac{(x-a)^r}{r!} b_r, \\ u_{n+1}(x) &= u_n(x) - L^{-1} \left[\sum_{j=0}^k \xi_j(x) u_n^{(j)}(x) - f(x) - \lambda_1 \int_a^x K_1(x,t) G_1(u_n(t)) dt - \lambda_2 \int_a^b K_2(x,t) G_2(u_n(t)) dt \right], \quad n \geq 0. \end{aligned} \tag{2.6}$$

The term $\sum_{r=0}^{k-1} \frac{(x-a)^r}{r!} b_r$ is obtained from the initial conditions, $\xi_k(x) \neq 0$. Relation (2.6) will enable us to determine the components $u_n(x)$ recursively for $n \geq 0$. Consequently, the approximation solution may be obtained by using $u(x) = \lim_{n \rightarrow \infty} u_n(x)$.

3. CONVERGENCE RESULTS

In this section, we shall give the convergence results of Eq. (1.1), with the initial condition (1.2) and prove it.

From Eq. (1.3), we assume:

$$L^{-1} \left[\int_a^x \frac{1}{\xi_k(x)} K_1(x, t) G_1(u_n(t)) dt \right] = \int_a^x \frac{(x-t)^k}{k! \xi_k(x)} K_1(x, t) G_1(u_n(t)) dt$$

and

$$\sum_{j=0}^{k-1} L^{-1} \left[\frac{\xi_j(x)}{\xi_k(x)} \right] u^{(j)}(x) = \sum_{j=0}^{k-1} \int_a^x \frac{(x-t)^{k-1} \xi_j(t)}{(k-1)! \xi_k(t)} u^{(j)}(t) dt.$$

We set,

$$\Psi(x) = L^{-1} \left[\frac{f(x)}{\xi_k(x)} \right] + \sum_{r=0}^{k-1} \frac{(x-a)^r}{r!} b_r.$$

Before starting and proving the main results, we introduce the following hypotheses:

(H1) There exist constants α, β and $\gamma_j > 0, j = 0, 1, \dots, k$ such that, for any $u_1, u_2 \in C(J, \mathbb{R})$

$$|G_1(u_1) - G_1(u_2)| \leq \alpha |u_1 - u_2|,$$

$$|G_2(u_1) - G_2(u_2)| \leq \beta |u_1 - u_2|,$$

$$|D^j(u_1) - D^j(u_2)| \leq \gamma_j |u_1 - u_2|,$$

we suppose that the nonlinear terms $G_1(u(x)), G_2(u(x))$ and $D^j(u) = (\frac{d^j}{dx^j})u(x) = \sum_{i=0}^{\infty} \gamma_{i,j}$, (D^j is a derivative operator), $j = 0, 1, \dots, k$, are Lipschitz continuous.

(H2) Suppose that for all $a \leq t \leq x \leq b$, and $j = 0, 1, \dots, k$:

$$\left| \frac{\lambda_1(x-t)^k K_1(x, t)}{k! \xi_k(x)} \right| \leq \theta_1, \quad \left| \frac{\lambda_1(x-t)^k K_1(x, t)}{k!} \right| \leq \theta_2,$$

$$\left| \frac{(x-t)^{k-1} \xi_j(t)}{(k-1)! \xi_k(t)} \right| \leq \theta_3, \quad \left| \frac{(x-t)^{k-1} \xi_j(t)}{(k-1)!} \right| \leq \theta_4,$$

$$\left| \lambda_2 L^{-1} \left[\frac{K_2(x, t)}{\xi_k(x)} \right] \right| \leq \theta_5, \quad \left| \lambda_2 L^{-1} [K_2(x, t)] \right| \leq \theta_6.$$

(H3) There exist three functions θ_3^*, θ_4^* , and $\gamma^* \in C(D, \mathbb{R}^+)$, the set of all positive function continuous on $D = \{(x, t) \in \mathbb{R} \times \mathbb{R} : 0 \leq t \leq x \leq 1\}$ such that:

$$\theta_3^* = \max |\theta_3|, \quad \theta_4^* = \max |\theta_4|, \quad \text{and} \quad \gamma^* = \max |\gamma_j|.$$

(H4) $\Psi(x)$ is bounded function for all x in $J = [a, b]$.

Theorem 3.1. *Suppose that (H1) – (H4), and if $0 < \psi < 1$, hold, the series solution $u(x) = \sum_{m=0}^{\infty} u_m(x)$ and $\|u_1\|_{\infty} < \infty$ obtained by the m -order deformation is convergent, then it converges to the exact solution of the problem (1.1) – (1.2).*

Proof. Denote as $(C[0, 1], \|\cdot\|)$ the Banach space of all continuous functions on J , with $|u_1(x)| < \infty$ for all x in J .

First we define the sequence of partial sums $\{s_n\}$, let s_n and s_m be arbitrary partial sums with $n \geq m$. We are going to prove that $s_n = \sum_{i=0}^n u_i(x)$ is a Cauchy sequence in this Banach space:

$$\begin{aligned} \|s_n - s_m\|_{\infty} &= \max_{\forall x \in J} |s_n - s_m| = \max_{\forall x \in J} \left| \sum_{i=0}^n u_i(x) - \sum_{i=0}^m u_i(x) \right| \\ &= \max_{\forall x \in J} \left| \sum_{i=m+1}^n u_i(x) \right| \\ &= \max_{\forall x \in J} \left| \sum_{i=m+1}^n \int_a^x \frac{\lambda_1(x-t)^k K_1(x,t)}{k! \xi_k(x)} A_{i-1} dt \right. \\ &\quad + \sum_{i=m+1}^n \int_a^b \lambda_2 L^{-1} \left[\frac{K_2(x,t)}{\xi_k(x)} \right] B_{i-1} dt \\ &\quad \left. - \sum_{j=0}^{k-1} \int_a^x \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} L_{(i-1)J} dt \right| \\ &= \max_{\forall x \in J} \left| \int_a^x \frac{\lambda_1(x-t)^k K_1(x,t)}{k! \xi_k(x)} \left(\sum_{i=m}^{n-1} A_i \right) dt \right. \\ &\quad + \int_a^b \lambda_2 L^{-1} \left[\frac{K_2(x,t)}{\xi_k(x)} \right] \left(\sum_{i=m}^{n-1} B_i \right) dt \\ &\quad \left. - \sum_{j=0}^{k-1} \int_a^x \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} \left(\sum_{i=m}^{n-1} L_{iJ} dt \right) \right|. \end{aligned}$$

From (2.2), we have

$$\begin{aligned} \sum_{i=m}^{n-1} A_i &= G_1(s_{n-1}) - G_1(s_{m-1}), \quad \sum_{i=m}^{n-1} B_i = G_2(s_{n-1}) - G_2(s_{m-1}), \\ \sum_{i=m}^{n-1} L_i &= D^j(s_{n-1}) - D^j(s_{m-1}). \end{aligned}$$

So,

$$\begin{aligned}
\|s_n - s_m\|_\infty &= \max_{x \in J} \left| \int_0^x \frac{\lambda_1(x-t)^k K_1(x,t)}{k! \xi_k(x)} [G_1(s_{n-1}) - G_1(s_{m-1})] dt \right. \\
&\quad + \int_a^b \lambda_2 L^{-1} \left[\frac{K_2(x,t)}{\xi_k(x)} \right] [G_2(s_{n-1}) - G_2(s_{m-1})] dt \\
&\quad \left. - \sum_{j=0}^{k-1} \int_a^x \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} [D^j(s_{n-1}) - D^j(s_{m-1})] dt \right| \\
&\leq \max_{x \in J} \int_0^x \left| \frac{\lambda_1(x-t)^k K_1(x,t)}{k! \xi_k(x)} \right| |G_1(s_{n-1}) - G_1(s_{m-1})| dt \\
&\quad + \int_a^b \left| \lambda_2 L^{-1} \left[\frac{K_2(x,t)}{\xi_k(x)} \right] \right| |G_2(s_{n-1}) - G_2(s_{m-1})| dt \\
&\quad + \sum_{j=0}^{k-1} \int_a^x \left| \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} \right| |D^j(s_{n-1}) - D^j(s_{m-1})| dt.
\end{aligned}$$

Let $n = m + 1$. Then

$$\begin{aligned}
\|s_n - s_m\|_\infty &\leq \psi \|s_m - s_{m-1}\|_\infty \\
&\leq \psi^2 \|s_{m-1} - s_{m-2}\|_\infty \\
&\quad \vdots \\
&\leq \psi^m \|s_1 - s_0\|_\infty,
\end{aligned}$$

so,

$$\begin{aligned}
\|s_n - s_m\|_\infty &\leq \|s_{m+1} - s_m\|_\infty + \|s_{m+2} - s_{m+1}\|_\infty + \cdots + \|s_n - s_{n-1}\|_\infty \\
&\leq [\psi^m + \psi^{m+1} + \cdots + \psi^{n-1}] \|s_1 - s_0\|_\infty \\
&\leq \psi^m [1 + \psi + \psi^2 + \cdots + \psi^{n-m-1}] \|s_1 - s_0\|_\infty \\
&\leq \psi^m \left(\frac{1 - \psi^{n-m}}{1 - \psi} \right) \|u_1\|_\infty.
\end{aligned}$$

Since $0 < \psi < 1$, we have $(1 - \psi^{n-m}) < 1$, then

$$\|s_n - s_m\|_\infty \leq \frac{\psi^m}{1 - \psi} \|u_1\|_\infty.$$

But $|u_1(x)| < \infty$, so, as $m \rightarrow \infty$, then $\|s_n - s_m\|_\infty \rightarrow 0$.

We conclude that $\{s_n\}$ is a Cauchy sequence in $C[0, 1]$, therefore $u = \lim_{n \rightarrow \infty} u_n$. Then, the series is convergence and the proof is complete. \square

Theorem 3.2. *If problem (1.1)–(1.2) has a unique solution, then the solution $u_n(x)$ obtained from the recursive relation (2.6) using VIM converges when $0 < \phi = (\alpha\theta_5 + \beta\theta_6 + k\gamma^*\theta_4^*)(b - a) < 1$.*

Proof. We have from equation (2.6):

$$\begin{aligned} u_{n+1}(x) - u(x) &= u_n(x) - u(x) - \left(L^{-1} \left[\sum_{j=0}^k \xi_j(x) [u_n^{(j)}(x) - u^{(j)}(x)] \right] \right. \\ &\quad - L^{-1} \left[\lambda_1 \int_a^x K_1(x, t) [G_1(u_n(t)) - G_1(u(t))] dt \right. \\ &\quad \left. \left. - L^{-1} \left[\lambda_2 \int_a^b K_2(x, t) [G_2(u_n(t)) - G_2(u(t))] dt \right] \right] \right). \end{aligned}$$

If we set, $\xi_k(x) = 1$, and $W_{n+1}(x) = u_{n+1}(x) - u(x)$, $W_n(x) = u_n(x) - u(x)$ since $W_n(a) = 0$, then

$$\begin{aligned} W_{n+1}(x) &= W_n(x) + \int_a^x \frac{\lambda_1 K_1(x, t)(x - t)^k}{k!} [G_1(u_n(t)) - G_1(u(t))] dt \\ &\quad + \int_a^b \lambda_2 L^{-1} [K_2(x, t) [G_2(u_n(t)) - G_2(u(t))] dt] - \sum_{j=0}^{k-1} \int_a^x \\ &\quad \times \frac{\xi_j(t)(x - t)^{k-1}}{(k - 1)!} [D^j(u_n(t)) - D^j(u(t))] dt - (W_n(x) - W_n(a)). \end{aligned}$$

Therefore,

$$\begin{aligned} |W_{n+1}(x)| &\leq \int_a^x \left| \frac{\lambda_1 K_1(x, t)(x - t)^k}{k!} \right| |W_n| \alpha dt \\ &\quad + \int_a^b \left| \lambda_2 L^{-1} [K_2(x, t)] \right| |W_n| \beta dt \\ &\quad + \sum_{j=0}^{k-1} \int_a^x \left| \frac{\xi_j(t)(x - t)^{k-1}}{(k - 1)!} \right| \max |\gamma_j| |W_n| dt \\ &\leq |W_n| \left[\int_a^x \alpha \theta_5 dt + \int_a^b \beta \theta_6 dt + \sum_{j=0}^{k-1} \int_a^x \theta_4^* \max |\gamma_j| \right] \\ &\leq |W_n| (\alpha\theta_5 + \beta\theta_6 + k\gamma^*\theta_4^*)(b - a) = |W_n| \phi. \end{aligned}$$

Hence,

$$\|W_{n+1}\| = \max_{\forall x \in J} |W_{n+1}(x)| \leq \phi \max_{\forall x \in J} |W_n(x)| = \phi \|W_n\|.$$

Since $0 < \phi < 1$, $\|W_n\| \rightarrow 0$. So, the series converges and the proof is complete. \square

4. APPLICATIONS

Example 4.1. Consider the following Volterra-Fredholm integro-differential equation.

$$u'(x) + xu(x) = 2x + x^3 - \frac{x^5}{5} - \frac{0.9^7}{7}x + \int_0^x u^2(t)dt + \int_0^{0.9} xu^3(t)dt,$$

with the initial condition $u(0) = u'(0) = 0$ and the the exact solution is $u(x) = x^2$.

TABLE 1. Numerical Results of the Example 4.1.

x	Exact solution	MDM	VIM
0.1	0.010000	0.016377	0.010024
0.2	0.040000	0.046990	0.040394
0.3	0.090000	0.094713	0.091969
0.4	0.160000	0.148751	0.151274
0.5	0.250000	0.236624	0.243752
0.6	0.360000	0.342563	0.350874
0.7	0.490000	0.478846	0.483681
0.8	0.640000	0.635372	0.630257
0.9	0.810000	0.790145	0.801487




FIGURE 1. Comparison Between MADM and VIM, for Example 4.1.

Example 4.2. Consider the nonlinear Volterra-Fredholm integro-differential equation:

$$(x^3 - 1)u^{(4)}(x) + (x^2 + 1)u''(x) = e^{-x}(x^2 + x^3) - \frac{x^2}{2e^2} - (0.130639)x + \int_0^x [u(t)]^2 dt + \int_0^{0.5} xt(1 + u^2(t))dt,$$

with the initial conditions: $u(0) = u''(0) = 1$, $u'(0) = u'''(0) = -1$, The exact solution is $u(x) = e^{-x}$.

TABLE 2. Numerical Results of the Example 4.2.

x	Exact solution	MADM	VIM
0.05	0.951229	0.908237	0.948796
0.1	0.904837	0.921381	0.908269
0.2	0.818731	0.796469	0.815826
0.3	0.740818	0.708649	0.739765
0.4	0.670320	0.649382	0.663792
0.5	0.606531	0.579846	0.607912




FIGURE 2. Comparison Between MADM and VIM, for Example 4.2.

5. CONCLUDING REMARKS

In this work, we have examined a few recent familiar numerical methods for solving integro-differential equations. The numerical studies showed that all the methods give highly accurate solutions for given equations. The MADM and the VIM are uncomplicated and comfortable. Despite this, they are not converging to a closed form. One advantage of VIM is that the initial solution can be freely chosen with some unknown parameters. An interesting point about this method is that with few number of iterations. The comparison reveals that although the numerical results of these methods are similar approximately, VIM is the easiest, the most efficient and convenient.

The problem considered in this paper can be generalized to a higher dimension involving a general formulation of fractional derivative with respect to another function. Also, study nonlinear fractional systems of Volterra-Fredholm integro-differential equations with nonlocal conditions is a direction which we are working on.

REFERENCES

- [1] S. Abbasbandy and S. Elyas, *Application of variational iteration method for system of nonlinear Volterra integro-differential equations*, Math. Comput. Appl. **2**(14) (2009), 147-158.
- [2] G. Adomian, *A review of the decomposition method in applied mathematics*, J. Math. Anal. Appl., **135**(2) (1988), 501-544.
- [3] S. Alao, F. Akinboro1, F. Akinpelu and R. Oderinu, *Numerical solution of integro-differential equation using Adomian decomposition and variational iteration methods*, IOSR J. Math., **10**(4) (2014), 18-22.
- [4] M.A. Araghi and S.S. Behzadi, *Solving nonlinear Volterra-Fredholm integro-differential equations using the modified Adomian decomposition method*, Comput. Methods Appl. Math., **9** (2009), 321-331.
- [5] S. Behzadi, S. Abbasbandy, T. Allahviranloo and A. Yildirim, *Application of homotopy analysis method for solving a class of nonlinear Volterra-Fredholm integro-differential equations*, J. Appl. Anal. Comput., **2**(2) (2012), 127-136.
- [6] M. Bani Issa and A. Hamoud, *Solving systems of Volterra integro-differential equations by using semi-analytical techniques*, Technology Reports of Kansai University, **62**(3) (2020), 685-690.
- [7] F.S. Fadhel, A.O. Mezaal and S.H. Salih, *Approximate solution of the linear mixed Volterra-Fredholm integro-differential equations of second kind by using variational iteration method*, Al- Mustansiriyah J. Sci., **24**(5) (2013), 137-146.
- [8] M.A. Fariborzi Araghi and S.S. Behzadi, *Solving nonlinear Volterra-Fredholm integral differential equations using the modified Adomian decomposition method*, Comput. Methods Appl. Math., **9** (2009), 1-11.
- [9] M. Ghasemi, M. kajani and E. Babolian, *Application of He's Homotopy Perturbation Method to Nonlinear Integro differential Equations*, Appl. Math. Comput., **188** (2007), 538-548.
- [10] J.H. He, *Variational iteration method for autonomous ordinary differential system*, Appl. Math. Comput., **114** (2000), 115-123.

- [11] J.H. He, *A variational approach to nonlinear problems and its application*, Mech. Appl., **20**(1) (1998), 30-34.
- [12] J.H. He and S.Q. Wang, *Variational iteration method for solving integro-differential equations*, Phys. Lett. A, **367** (2007), 188-191.
- [13] A. Hamoud and K. Ghadle, *Some new existence, uniqueness and convergence results for fractional Volterra-Fredholm integro-differential equations*, J. Appl. Comput. Mech., **5**(1) (2019), 58-69.
- [14] A. Hamoud and K. Ghadle, *The approximate solutions of fractional Volterra-Fredholm integro-differential equations by using analytical techniques*, Probl. Anal. Issues Anal., **7**(25) (2018), 41-58.
- [15] A. Hamoud, M. SH. Bani Issa and K. Ghadle, *Existence and uniqueness results for nonlinear Volterra-Fredholm integro-differential equations*, Nonlinear Funct. Anal. Appl., **23**(4) (2018), 797-805
- [16] A. Hamoud and K. Ghadle, *Homotopy analysis method for the first order fuzzy Volterra-Fredholm integro-differential equations*, Indonesian J.f Electrical Eng. Comput. Sci., **11**(3) (2018), 857-867.
- [17] A. Hamoud and K. Ghadle, *Modified Adomian decomposition method for solving fuzzy Volterra-Fredholm integral equation*, J. Indian Math. Soc., **85**(1-2) (2018), 53-69.
- [18] A. Hamoud, K. Ghadle and S. Atshan, *The approximate solutions of fractional integro-differential equations by using modified Adomian decomposition method*, Khayyam J. Math., **5**(1) (2019), 21-39.
- [19] A. Hamoud and K. Ghadle, *Modified Laplace decomposition method for fractional Volterra-Fredholm integro-differential equations*, J. Math Mdel., **6**(1) (2018), 91-104.
- [20] A. Hamoud, A. Azeez and K. Ghadle, *A study of some iterative methods for solving fuzzy Volterra-Fredholm integral equations*, Indonesian J. Electrical Eng. Comput. Sci., **11**(3) (2018), 1228-1235.
- [21] A. Hamoud, K. Hussain and K. Ghadle, *The reliable modified Laplace Adomian decomposition method to solve fractional Volterra-Fredholm integro differential equations*, Dyn. Continuous, Disc. Impul. Sys. Series B: Appl. Algo., **26** (2019), 171-184.
- [22] K. Hussain, A. Hamoud and N. Mohammed, *Some new uniqueness results for fractional integro-differential equations*, Nonlinear Funct. Anal. Appl., **24**(4) (2019), 827-836.
- [23] H. Han, Y. Liu, T. Lu and X. He, *New algorithm for the system of nonlinear weakly singular Volterra integral equations of the second kind and integro-differential equations*, J. Inform. Comput. Sci. **7** (2010), 1229-1235.
- [24] S.A. Khuri, *A Laplace decomposition algorithm applied to class of nonlinear differential equations*, J. Math. Appl. **4** (2001), 141-155.
- [25] R. Mittal and R. Nigam, *Solution of fractional integro-differential equations by Adomian decomposition method*, Int. J. Appl. Math. Mech. **4**(2) (2008), 87-94.
- [26] N. Ngarhasta, B. Some, K. Abbaoui, and Y. Cherruau, *New numerical study of Adomian method applied to a diffusion model*, Kybernetes, **31** (2002), 61-75.
- [27] S.B. Shadan, *The use of iterative method to solve two-dimensional nonlinear Volterra-Fredholm integro-differential equations*, J. Comm. Numer. Anal., (2012), 1-20.
- [28] Y. Salih and S. Mehmet, *The approximate solution of higher order linear Volterra-Fredholm integro-differential equations in term of Taylor polynomials*, Appl. Math. Comput., **112** (2000), 291-308.
- [29] A.M. Wazwaz, *The modified decomposition method for analytic treatment of nonlinear integral equations and systems of nonlinear integral equations*, Int. J. Compu. Math., **82**(9) (2005), 1107-1115.