



## DEGREE OF APPROXIMATION OF A FUNCTION ASSOCIATED WITH HARDY-LITTLEWOOD SERIES IN WEIGHTED ZYGMUND $W(Z_r^{(\omega)})$ -CLASS USING EULER-HAUSDORFF SUMMABILITY MEANS

Tejaswini Pradhan<sup>1</sup> and G. V. V. Jagannadha Rao<sup>2</sup>

<sup>1</sup>Department of Mathematics, Kalinga University, Naya Raipur,  
Chhattisgarh - 492101, India  
e-mail: [tejaswini.pradhan@kalingauniversity.ac.in](mailto:tejaswini.pradhan@kalingauniversity.ac.in)

<sup>2</sup>Department of Mathematics, Kalinga University, Naya Raipur,  
Chhattisgarh - 492101, India  
e-mail: [dr.gvvj.rao@kalingauniversity.ac.in](mailto:dr.gvvj.rao@kalingauniversity.ac.in)

**Abstract.** Approximation of functions of Lipschitz and Zygmund classes have been considered by various researchers under different summability means. In the proposed study, we investigated an estimation of the order of convergence of a function associated with Hardy-Littlewood series in the weighted Zygmund class  $W(Z_r^{(\omega)})$ -class by applying Euler-Hausdorff summability means and subsequently established some (presumably new) results. Moreover, the results obtained here represent the generalization of several known results.

### 1. INTRODUCTION

Summability methods have been used in various fields of mathematics. For example, summability methods are applied in function theory in connection with the analytic continuation of holomorphic functions and the boundary behavior of a power series, in applied analysis for generation of iteration methods for finding solutions of a system of equations, and for acceleration of convergence in approximation theory. Also, it has been used in other fields

---

<sup>0</sup>Received April 19, 2023. Revised June 22, 2023. Accepted June 27, 2023.

<sup>0</sup>2020 Mathematics Subject Classification: 41A24, 41A25, 42B05, 42B08.

<sup>0</sup>Keywords: Degree of approximation, weighted Zygmund class, Euler-Hausdorff mean, Hardy-Littlewood series.

<sup>0</sup>Corresponding author: G. V. V. Jagannadha Rao

([dr.gvvj.rao@kalingauniversity.ac.in](mailto:dr.gvvj.rao@kalingauniversity.ac.in)).

of mathematics like probability theory (Markov chains) and number theory (Prime number theorem). The approximation analysis of signals (functions) has great importance in the field of science and engineering. It has also given a new aspect due to its exhaustive appliance in signal analysis, system design, modern telecommunications, radar and image processing system. The error estimation of functions in various function spaces such as Lipschitz, Hölder, Zygmund, Besov spaces using different summability techniques of Fourier series has been received a growing interest of several researchers in the last decades. Functions in  $L_r$  ( $r \geq 1$ )-spaces assumed to be most practicable in signal analysis. Particularly,  $L_1$ ,  $L_2$  and  $L_\infty$  spaces are used by engineers for designing digital filters. The generalized Zygmund class  $Z_r^{(\omega)}$  ( $r \geq 1$ ) is a generalization of  $Z_{(\alpha)}$ ,  $Z_{(\alpha),r}$ ,  $Z^{(\omega)}$ -class.

The generalized Zygmund class  $Z_r^{(\omega)}$  ( $r \geq 1$ ) is investigated by Leindler [7], Moricz [9], Moricz and Nemeth [10]. Lal and Shireen [6] established results on approximation of functions of generalized Zygmund class by Matrix-Euler summability mean of Fourier series. Pradhan *et al.* [14] studied on approximation of signals belonging to generalized Lipschitz class using  $(\bar{N}, p_n, q_n)(E, s)$ -summability mean of Fourier series. Singh *et al.* [16] studied approximation of functions in the generalized Zygmund class using Hausdorff means. Pradhan *et al.* [13] studied on approximation of signals in the generalized Zygmund class via  $(E, 1)(\bar{N}, p_n)$  summability means of conjugate Fourier series. In 2019, Pradhan *et al.* [15] studied approximation of signals using generalized Zygmund class using  $(E, 1)(\bar{N}, p_n)$  summability means of Fourier series. Das *et al.* [1] proved approximation of functions in the weighted Zygmund class via Euler-Hausdorff product summability means of Fourier series. Again, in 2020, Padhy *et al.* [12] estimated the degree of approximation of functions of generalized Zygmund class associated with Hardy-Littlewood series using Riesz mean. Very recently in 2023, Jena *et al.* [4] studied on the degree of approximation of Fourier series based on a certain class of product deferred summability means.

Motivated by the above mentioned works, to get best approximation and advance study in this direction, In this proposed paper, we give an estimation of degree of approximation of functions associated with Hardy-Littlewood series in weighted Zygmund class using Euler-Hausdorff summability means.

## 2. PRELIMINARIES

Let  $f(x)$  be a periodic function of period  $2\pi$ , which is Lebesgue integrable in  $[-\pi, \pi]$  and Fourier series associated with  $f(x)$  is given by

$$\sum_{n=0}^n A_n(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx). \quad (2.1)$$

Let  $S_n^M(x)$  denotes the  $n^{\text{th}}$  partial sum of the (2.1) is given by

$$S_n^M(x) = \sum_{k=0}^{n-1} A_k(x) + \frac{A_n(x)}{2}. \quad (2.2)$$

Then the Hardy-Littlewood series or HL-series associated with  $f(x)$  is given by

$$\frac{c_0}{2} + \sum_{n=1}^{\infty} \frac{S_n^M(x) - f(x)}{n}, \quad (2.3)$$

where

$$c_0 = \frac{2}{\pi} \int_0^{\pi} \phi(x, u) \frac{u}{2} \cot \frac{u}{2} du$$

and  $\phi(x, u) = f(x+u) + f(x-u) - 2f(x)$ . Let

$$\eta(u) = \int_u^{\pi} \phi(x, u) \frac{1}{2} \cot \frac{u}{2} du. \quad (2.4)$$

Clearly,  $\eta(u)$  is an even function and Lebesgue integrable in  $[-\pi, \pi]$ . Also, the HL-series (2.3) is the Fourier series of  $\eta(u)$  at  $u = 0$ .

Let us write  $\xi_n(f, x) = \frac{2}{\pi} \int_0^{\pi} \eta(u) \frac{\sin(n+\frac{1}{2})u}{2\sin\frac{u}{2}} du$ , which represent the  $n^{\text{th}}$  partial sum of  $\eta(u)$ .

The  $L_r$  norm of a function  $\eta$  is defined by

$$\|\eta\|_r = \begin{cases} \left( \frac{1}{2\pi} \int_0^{2\pi} |\eta(x)|^r dx \right)^{\frac{1}{r}}, & 1 \leq r < \infty, \\ \text{ess sup}_{0 < x \leq 2\pi} |\eta(x)|, & r = \infty. \end{cases}$$

The degree of approximation of a function  $\eta : \mathbb{R} \rightarrow \mathbb{R}$  by a trigonometric polynomial  $t_n$  of order  $n$  under  $\|\cdot\|_{L_\infty}$  norm is defined as

$$\|t_n - \eta(x)\|_{L_\infty} = \sup_{x \in \mathbb{R}} |t_n(x) - \eta(x)|$$

and let a function  $\eta \in L_r$ , its degree of approximation  $E_n(\eta)$  is given by

$$E_n(\eta) = \min_{t_n} \|t_n - \eta\|_{L_r}.$$

Zygmund modulus of continuity [18] of  $\eta$  is defined by

$$\omega(\eta, h) = \sup_{0 \leq h, x \in \mathbb{R}} |\eta(x+h) + \eta(x-h) - 2\eta(x)|.$$

Let  $C_{2\pi}$  denote the Banach space of all  $2\pi$ -periodic continuous functions defined on  $[0, 2\pi]$  under the supremum norm. For  $0 < \alpha \leq 1$ , the function space

$$Z_{(\alpha)} = \{ \eta \in C_{2\pi} : |\eta(x+t) + \eta(x-t) - 2\eta(x)| = O(|t|^\alpha) \}$$

is a Banach space under the norm  $\| \cdot \|_{(\alpha)}$  is defined by

$$\| \eta \|_{(\alpha)} = \sup_{0 \leq x \leq 2\pi} |\eta(x)| + \sup_{x, t \neq 0} \frac{|\eta(x+t) + \eta(x-t) - 2\eta(x)|}{|t|^\alpha}.$$

For  $\eta \in L_r[0, 2\pi], r \geq 1$ , the integral Zygmund modulus of continuity is defined by

$$\omega_r(\eta, h) = \sup_{0 < t \leq h} \left\{ \frac{1}{2\pi} \int_0^{2\pi} |\eta(x+t) + \eta(x-t) - 2\eta(x)|^r dx \right\}^{\frac{1}{r}}.$$

Moreover, for  $\eta \in C_{2\pi}$  and  $r = \infty$ ,

$$\omega_\infty(\eta, h) = \sup_{0 < t \leq h} \max_x |\eta(x+t) + \eta(x-t) - 2\eta(x)|.$$

Also, it is known that  $\omega_r(\eta, h) \rightarrow 0$  as  $r \rightarrow 0$ .

We now define,

$$Z_{(\alpha),r} = \left\{ \eta \in L_r[0, 2\pi] : \left( \int_0^{2\pi} |\eta(x+t) + \eta(x-t) - 2\eta(x)|^r dx \right)^{\frac{1}{r}} = O(|t|^\alpha) \right\}.$$

The space  $Z_{(\alpha),r}, r \geq 1, 0 < \alpha \leq 1$  is a Banach space under the norm  $\| \cdot \|_{(\alpha),r}$  and that,

$$\| \eta \|_{(\alpha),r} = \| \eta \|_r + \sup_{t \neq 0} \frac{\| \eta(\cdot + t) + \eta(\cdot - t) - 2\eta(\cdot) \|_r}{|t|^\alpha}.$$

The class of function  $Z^{(\omega)}$  is defined as

$$Z^{(\omega)} = \{ \eta \in C_{2\pi} : |\eta(x+t) + \eta(x-t) - 2\eta(x)| = O(\omega(t)) \},$$

where  $\omega$  is a Zygmund modulus of continuity, that is,  $\omega$  is positive, non-decreasing continuous function with the sub linearity property, that is,

- (i)  $\omega(0) = 0$ ,
- (ii)  $\omega(t_1 + t_2) \leq \omega(t_1) + \omega(t_2)$ .

Let  $\omega : [0, 2\pi] \rightarrow \mathbb{R}$  be an arbitrary function with  $\omega(t) > 0$  for  $0 \leq t < 2\pi$  and let  $\lim_{t \rightarrow 0^+} \omega(t) = \omega(0) = 0$ , define

$$Z_r^{(\omega)} = \left\{ \eta \in L_r : 1 \leq r \leq \infty, \sup_{t \neq 0} \frac{\| \eta(\cdot + t) + \eta(\cdot - t) - 2\eta(\cdot) \|_r}{\omega(t)} < \infty \right\},$$

where

$$\|\eta\|_r^{(\omega)} = \|\eta\|_r + \sup_{t \neq 0} \frac{\|\eta(\cdot + t) + \eta(\cdot - t) - 2\eta(\cdot)\|_r}{\omega(t)}, \quad r \geq 1.$$

Then, clearly  $\|\cdot\|_r^{(\omega)}$  is a norm on  $Z_r^{(\omega)}$ . As we know  $L_r$  ( $r \geq 1$ ) is complete, the space  $Z_r^{(\omega)}$  is also complete. Hence we can say  $Z_r^{(\omega)}$  is a Banach space under the norm  $\|\cdot\|_r^{(\omega)}$ .

Now we define the weighted Zygmund class as

$$\begin{aligned} & W(Z_r^{(\omega)}) \\ &= \left\{ \eta \in W(Z_r^{(\omega)}) : 1 \leq r \leq \infty, \sup_{t \neq 0} \frac{\|(\eta(\cdot + t) + \eta(\cdot - t) - 2\eta(\cdot)) \sin^\beta(\cdot)\|_r}{\omega(t)} \leq \infty \right\}, \end{aligned} \quad (2.5)$$

where

$$\|\eta\|_r^{(\omega)} = \|\eta\|_r + \sup_{t \neq 0} \frac{\|(\eta(\cdot + t) + \eta(\cdot - t) - 2\eta(\cdot)) \sin^\beta(\cdot)\|_r}{\omega(t)}, \quad r \geq 1. \quad (2.6)$$

Clearly,  $\|\cdot\|_r^{*(\omega)}$  is a norm of  $Z_r^{(\omega)}$ . The space  $Z_r^{(\omega)}$  is complete because  $L_r$ ,  $r \geq 1$  is complete. Hence, we can say that  $W(Z_r^{(\omega)})$  is complete.

As  $Z_r^{(\omega)}$  is a Banach space under  $\|\cdot\|_r^{(\omega)}$ , so  $W(Z_r^{(\omega)})$  is also a Banach space under  $\|\cdot\|_r^{(\omega)}$  norm.

- (i) If we put  $\beta = 0$  in  $W(Z_r^{(\omega)})$  class, then it reduces to  $Z_r^{(\omega)}$  class.
- (ii) If we put  $r \rightarrow \infty$ , then the class  $Z_r^{(\omega)}$  reduces to the  $Z^{(\omega)}$  class.
- (iii) If we put  $\omega(t) = t^\alpha$  in  $Z_r^{(\omega)}$  class, then it reduces to  $Z_{(\alpha),r}$  class.
- (iv) If we put  $\omega(t) = t^\alpha$ , the  $Z^{(\omega)}$  class reduces to  $Z_{(\alpha)}$  class.

Here  $\omega(t)$  and  $v(t)$  denotes the Zygmund moduli of continuity such that  $\left(\frac{\omega(t)}{v(t)}\right)$  is positive, non-decreasing, then

$$\|\eta\|_r^{(v)} \leq \max\left(1, \frac{\omega(2\pi)}{v(2\pi)}\right) \|\eta\|_r^{(\omega)} \leq \infty.$$

Thus, we have

$$Z_r^{(\omega)} \subseteq Z_r^{(v)} \subseteq L_r \quad (r \geq 1).$$

Hence,

$$W(Z_r^{(\omega)}) \subseteq W(Z_r^{(v)}) \subseteq W(L_r, \omega(t)).$$

Hausdorff matrices were first introduced by Hurwitz and Silverman [3] as the collection of lower triangular matrices that commute the Cesaro matrix of

order one. A Hausdorff matrix  $H \equiv (h_{n,k})$  is an infinite lower matrix defined by

$$h_{n,k} = \begin{cases} \binom{n}{k} \Delta^{n-k} \mu_k, & \text{if } 0 \leq k \leq n, \\ 0, & \text{if } k > n, \end{cases}$$

where the operator  $\Delta$  is defined by  $\Delta \mu_n \equiv \mu_n - \mu_{n+1}$  and  $\Delta^{k+1} \mu_n \equiv \Delta^k(\Delta \mu_n)$ .

Let  $\sum_{n=0}^\infty u_n$  be an infinite series with partial sum  $s_n = \sum_{k=0}^n u_k$ . If  $t_n^H = \sum_{k=0}^n h_{n,k} s_k \rightarrow s$  as  $n \rightarrow \infty$ ,  $\sum_{k=0}^\infty u_n$  is said to be summable to  $s$  by the Hausdorff matrix summability method ( $\Delta_H$  means). The Hausdorff matrix  $H$  is regular, that is,  $H$  preserves the limit of each convergent sequence if and only if

$$\int_0^1 |d(\alpha(z))| < \infty,$$

where the mass function  $\alpha \in BV[0, 1]$ ,  $\alpha(0+) = \alpha(0) = 0$  and  $\alpha(1) = 1$ . In this case, the  $\mu_n$  has the representation

$$\mu_n = \int_0^1 z^n d\alpha z.$$

Let  $E_n^{(q)} = \frac{1}{(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} s_k$ ,  $q > 0$ . If  $E_n^{(q)} \rightarrow s$  as  $n \rightarrow \infty$ ,  $\sum_{k=0}^\infty u_n$  is said to be summable to  $s$  by Euler method, that is, the  $(E, q)$  method (see [1]). The  $(E, q)$  transform of  $t_n^H$  transform defines the  $E^{(q)} \cdot \Delta_H$  transform of  $(s_n)$ . It is denoted by  $T_n^{EH}$ . Thus,

$$T_n^{EH} = \frac{1}{(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sum_{v=0}^k h_{k,v} s_v.$$

If  $T_n^{EH} \rightarrow s$  as  $n \rightarrow \infty$ ,  $\sum_{k=0}^\infty u_n$  is said to be summable to  $s$  by Euler-Hausdorff summability means (see [5]), that is, the  $E^{(q)} \cdot \Delta_H$  means. As the Euler method and Hausdorff methods are regular,  $E^{(q)} \cdot \Delta_H$  method is regular.

We use the following notations throughout the papers,

$$\phi(x, t) = f(x + t) + f(x - t) - f(x),$$

$$\begin{aligned} K_n^{EH} &= \frac{1}{2\pi(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sum_{v=0}^k h_{k,v} \frac{\sin(v + \frac{1}{2})t}{\sin(\frac{t}{2})} \\ &= \frac{1}{2\pi(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sum_{v=0}^k \int_0^1 z^k (1-z)^{k-v} d\alpha(z) \frac{\sin(v + \frac{1}{2})t}{\sin(\frac{t}{2})}. \end{aligned} \tag{2.7}$$

## 3. MAIN RESULTS

To prove the main results, we need the followings lemmas.

**Lemma 3.1.**  $|K_n^{EH}(t)| = O(n+1)$  for  $0 \leq t \leq \frac{1}{n+1}$ .

*Proof.* For  $\sin nt \leq n \sin t$ , we have

$$\begin{aligned} |K_n^{EH}(t)| &= \left| \frac{1}{2\pi(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sum_{v=0}^k \int_0^1 z^k (1-z)^{k-v} d\alpha(z) \frac{\sin(v + \frac{1}{2})t}{\sin(\frac{t}{2})} \right| \\ &\leq \frac{1}{2\pi(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sum_{v=0}^k \int_0^1 z^k (1-z)^{k-v} d\alpha(z) \left| \frac{\sin(v + \frac{1}{2})t}{\sin(\frac{t}{2})} \right| \\ &\leq \frac{1}{2\pi(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sum_{v=0}^k \int_0^1 z^k (1-z)^{k-v} d\alpha(z) \frac{(2v+1) \sin \frac{1}{2}t}{\sin(\frac{t}{2})} \\ &= \frac{(2n+1)N}{2\pi(1+q)^n} \binom{n}{k} q^{n-k} \sum_{v=0}^k \int_0^1 z^k (1-z)^{k-v} dz \\ &= O(n+1), \end{aligned} \tag{3.1}$$

where  $N = \sup_{0 < z \leq 1} |\alpha'(z)|$ .  $\square$

**Lemma 3.2.**  $|K_n^{EH}(t)| = O\left(\frac{1}{(n+1)t^2}\right)$  for  $\frac{1}{n+1} \leq t \leq \pi$ .

*Proof.* For  $|\sin nt| = 1$  and  $\sin \frac{t}{2} \geq \frac{t}{\pi}$ . First we calculate,

$$\begin{aligned} &\int_0^1 \sum_{k=0}^n \binom{n}{k} z^k (1-z)^{n-k} \sin(n + \frac{1}{2})t \, d\alpha(z) \\ &= \int_0^1 \operatorname{Im} \left[ \sum_{k=0}^n \binom{n}{k} z^k (1-z)^{n-k} e^{i(n+\frac{1}{2})t} \, d\alpha(z) \right] \\ &= \int_0^1 \operatorname{Im} \left[ e^{i\frac{t}{2}} (1-z)^n \sum_{k=0}^n \binom{n}{k} \left(\frac{z}{1-z}\right)^k e^{ikt} \, d\alpha(z) \right] \\ &= \int_0^1 \operatorname{Im} \left[ e^{i\frac{t}{2}} (1-z + ze^{it})^n \, d\alpha(z) \right] \\ &= N \operatorname{Im} \left( \frac{(1-z + ze^{it})^{n+1}}{e^{i\frac{t}{2}} - e^{-i\frac{t}{2}}(n+1)} \right)_{z=0}^{z=1} \\ &= N \operatorname{Im} \left( \frac{e^{i(n+1)t} - 1}{2i \sin(\frac{t}{2})(n+1)} \right) \end{aligned}$$

$$\begin{aligned}
 &= \frac{N}{2n+1} \operatorname{Im} \left( \frac{\cos(n+1)t + i \sin(n+1)t - 1}{2i \sin \frac{t}{2}} \right) \\
 &= \frac{N}{2n+1} \left( \frac{\sin(n+1)t}{i \sin \frac{t}{2}} \right) \\
 &= \frac{N}{2n+1} \times \frac{\pi}{t} \\
 &= O \left( \frac{1}{(n+1)t} \right). \tag{3.2}
 \end{aligned}$$

Now applying Jordan’s lemma, we have

$$\begin{aligned}
 |K_n^{EH}(t)| &= \left| \frac{1}{2\pi(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sum_{v=0}^k \int_0^1 z^k (1-z)^{k-v} d\alpha(z) \frac{\sin(v + \frac{1}{2})t}{\sin(\frac{t}{2})} \right| \\
 &\leq \frac{1}{2t(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \left| \sum_{v=0}^k \int_0^1 z^k (1-z)^{k-v} \sin(v + \frac{1}{2})t d\alpha(z) \right| \\
 &= \frac{1}{2t(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} O \left( \frac{1}{(k+1)t} \right) \\
 &= \left( \frac{1}{2t^2(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \left( \frac{1}{(k+1)} \right) \right) \\
 &= O \left( \frac{1}{(n+1)t^2} \right). \tag{3.3}
 \end{aligned}$$

□

**Lemma 3.3.** *Let  $f \in Z_r^{(\omega)}$ . Then for  $0 < t \leq \pi$ ,*

- (1)  $\|\phi(\cdot, t)\|_r = O(\omega(t))$ ,
- (2)  $\|\phi(\cdot + y, t) + \phi(\cdot - y, t) - 2\phi(\cdot, t)\|_r = \begin{cases} O(\omega(t)), \\ O(\omega(y)), \end{cases}$
- (3) *If  $\omega$  and  $v$  denotes the zygmond moduli such that  $(\frac{\omega(t)}{v(t)})$  is positive and increasing, then we have*

$$\|\phi(\cdot + y, t) + \phi(\cdot - y, t) - 2\phi(\cdot, t)\|_r = O \left( v(y) \frac{\omega(t)}{v(t)} \right),$$

where  $\phi(x, t) = f(x + t) + f(x - t) - 2f(x)$ .

**Lemma 3.4.**  $\|(\phi(\cdot + y, t) + \phi(\cdot - y, t) - 2\phi(\cdot, t)) \sin^\beta(\cdot)\|_r = O \left( t^\beta v(y) \left( \frac{\omega(t)}{v(t)} \right) \right)$ .



*Proof.* For  $v$  is positive, nondecreasing,  $t \leq y$ ,  $|\sin^\beta t| \leq t^\beta$  and using Lemma 3.3, we obtained

$$\begin{aligned} \|(\phi(\cdot + y, t) + \phi(\cdot - y, t) - 2\phi(\cdot, t)) \sin^\beta(\cdot)\|_r &= O(t^\beta \omega(t)) \\ &= O\left(t^\beta v(t) \left(\frac{\omega(t)}{v(t)}\right)\right) \\ &\leq O\left(t^\beta v(y) \left(\frac{\omega(t)}{v(t)}\right)\right). \end{aligned}$$

Since  $\frac{\omega(t)}{v(t)}$  is positive, non-decreasing, if  $t \geq y$ , then  $\frac{\omega(t)}{v(t)} \geq \frac{\omega(y)}{v(y)}$ , so that

$$\begin{aligned} \|(\phi(\cdot + y, t) + \phi(\cdot - y, t) - 2\phi(\cdot, t)) \sin^\beta(\cdot)\|_r &= O(t^\beta \omega(y)) \\ &= O\left(t^\beta v(y) \left(\frac{\omega(t)}{v(t)}\right)\right). \end{aligned}$$

□

The main objective of this paper is to prove the following theorems.

**Theorem 3.5.** *Let  $\eta$  be a  $2\pi$  periodic function and Lebesgue integrable on  $[-\pi, \pi]$  and belonging to weighted Zygmund class  $W(Z_r^{(\omega)})$ ,  $r \geq 1$ . Then the degree of approximation of signal (function)  $\eta$ , using Euler-Hausdorff summability means of HL-series (2.3) is given by*

$$E_n(\eta) = \inf \|T_n^{EH} - \eta\|_r = O\left(\frac{1}{(n+1)^2} \int_{\frac{1}{n+1}}^\pi \frac{t^{\beta-2} \omega(t)}{v(t)} dt\right), \quad (3.4)$$

where  $\omega$  and  $v$  denotes the zygmond moduli such that  $(\frac{\omega(t)}{v(t)})$  is positive and increasing.

*Proof.* Following the results of Titchmarsh [17], the Euler-Hausdorff transform of  $\{\xi_n(f, x)\}$  and is denoted by

$$\begin{aligned} \tau_n^{EH}(x) - \eta(x) &= \frac{2}{\pi} \int_0^\pi \phi(x, t) \frac{1}{(1+q)^n} \sum_{v=0}^n \binom{n}{v} q^{n-v} \{t_n^H(x) - \eta(x)\} \\ &= \frac{2}{\pi} \int_0^\pi \phi(x, t) \frac{1}{(1+q)^n} \sum_{v=0}^n \binom{n}{v} q^{n-v} \\ &\quad \times \left\{ \frac{1}{2\pi} \int_0^\pi \phi(x; t) \sum_{v=0}^k \int_0^1 \binom{k}{v} z^v (1-z)^{k-v} d\alpha(z) \frac{\sin(v + \frac{1}{2})t}{\sin \frac{t}{2}} dt \right\} \\ &= \int_0^\pi \phi(x; t) K_n^{EH}(t) dt \\ &= \mathcal{L}_n(x). \end{aligned} \quad (3.5)$$

Now

$$\mathcal{L}_n(x) = T_n^{EH}(x) - f(x) = \int_0^\pi \phi(x; t) K_n^{EH}(t) dt. \quad (3.6)$$

This implies

$$\begin{aligned} & \mathcal{L}_n(x+y) + \mathcal{L}_n(x-y) - 2\mathcal{L}_n(x) \\ &= \int_0^\pi [\phi(x+y; t) + \phi(x-y; t) - 2\phi(x; t)] K_n^{EH}(t) dt, \end{aligned} \quad (3.7)$$

$$\begin{aligned} & (\mathcal{L}_n(\cdot+y) + \mathcal{L}_n(\cdot-y) - 2\mathcal{L}_n(\cdot)) \sin^\beta(\cdot) \\ &= \int_0^\pi \left( (\phi(\cdot+y; t) + \phi(\cdot-y; t) - 2\phi(\cdot; t)) \sin^\beta(\cdot) \right) K_n^{EH}(t) dt. \end{aligned} \quad (3.8)$$

Now we can write

$$\begin{aligned} & \|(\mathcal{L}_n(\cdot+y) + \mathcal{L}_n(\cdot-y) - 2\mathcal{L}_n(\cdot)) \sin^\beta(\cdot)\|_r \\ &= \int_0^\pi \|(\phi(\cdot+y; t) + \phi(\cdot-y; t) - 2\phi(\cdot; t)) \sin^\beta(\cdot)\|_r K_n^{EH}(t) dt \\ &= \int_0^{\frac{1}{n+1}} \|(\phi(\cdot+y; t) + \phi(\cdot-y; t) - 2\phi(\cdot; t)) \sin^\beta(\cdot)\|_r K_n^{EH}(t) dt \\ &\quad + \int_{\frac{1}{n+1}}^\pi \|(\phi(\cdot+y; t) + \phi(\cdot-y; t) - 2\phi(\cdot; t)) \sin^\beta(\cdot)\|_r K_n^{EH}(t) dt \\ &:= I_1 + I_2. \end{aligned} \quad (3.9)$$

Further the function  $f \in W(Z_r^{(\omega)})$  implies  $\phi \in W(Z_r^{(\omega)})$  and applying Lemma 3.1, Lemma 3.4 and monotonicity of  $\frac{\omega(t)}{v(t)}$  with respect to  $t$ , we have

$$\begin{aligned} I_1 &= \int_0^{\frac{1}{n+1}} \|(\phi(\cdot+y; t) + \phi(\cdot-y; t) - 2\phi(\cdot; t)) \sin^\beta(\cdot)\|_r K_n^{EH}(t) dt \\ &= O\left(\int_0^{\frac{1}{n+1}} v(y) \frac{t^\beta \omega(t)}{v(t)} (n+1) dt\right) \\ &= O\left((n+1) v(y) \int_0^{\frac{1}{n+1}} \frac{t^\beta \omega(t)}{v(t)} dt\right) \\ &= O\left((n+1) v(y) \frac{\omega(\frac{1}{n+1})}{v(\frac{1}{n+1})} \int_0^{\frac{1}{n+1}} t^\beta dt\right) \\ &= O\left(\frac{1}{(n+1)^\beta} v(y) \frac{\omega(\frac{1}{n+1})}{v(\frac{1}{n+1})}\right). \end{aligned} \quad (3.10)$$

Next, using Lemma 3.2 and Lemma 3.4, we get

$$\begin{aligned} I_2 &= \int_{\frac{1}{n+1}}^{\pi} \|(\phi(\cdot + y; t) + \phi(\cdot - y; t) - 2\phi(\cdot; t)) \sin^{\beta}(\cdot)\|_r K_n^{EH}(t) dt \\ &= O\left(\int_{\frac{1}{n+1}}^{\pi} v(y) \frac{t^{\beta} \omega(t)}{v(t)} \frac{1}{(n+1)t^2} dt\right) \\ &= O\left(\frac{1}{(n+1)} v(y) \int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-2} \omega(t)}{v(t)} dt\right). \end{aligned} \quad (3.11)$$

Thus using (3.9), (3.10) and (3.11) we can write

$$\begin{aligned} &\|(\mathcal{L}_n(\cdot + y) + \mathcal{L}_n(\cdot - y) - 2\mathcal{L}_n(\cdot)) \sin^{\beta}(\cdot)\|_r \\ &= O\left(\frac{1}{(n+1)^{\beta}} v(y) \frac{\omega(\frac{1}{n+1})}{v(\frac{1}{n+1})}\right) + O\left(\frac{1}{(n+1)} v(y) \int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-2} \omega(t)}{v(t)} dt\right). \end{aligned} \quad (3.12)$$

Therefore, we have

$$\begin{aligned} &\sup_{y \neq 0} \frac{\|\mathcal{L}_n(\cdot + y) + \mathcal{L}_n(\cdot - y) - 2\mathcal{L}_n(\cdot)\|_r}{v(y)} \\ &= O\left(\frac{1}{(n+1)^{\beta}} \frac{\omega(\frac{1}{n+1})}{v(\frac{1}{n+1})}\right) + O\left(\frac{1}{(n+1)} \int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-2} \omega(t)}{v(t)} dt\right). \end{aligned} \quad (3.13)$$

Clearly,

$$\phi(x; t) = |f(x+t) + f(x-t) - 2f(x)|.$$

Now applying Minkowski's inequality, we have

$$\|\phi(x; t)\|_r = \|f(x+t) + f(x-t) - 2f(x)\|_r. \quad (3.14)$$

Now using Lemma 3.3, we have

$$\begin{aligned} &\|(\mathcal{L}_n(\cdot)) \sin^{\beta}(\cdot)\|_r \\ &\leq \left(\int_0^{\frac{1}{n+1}} + \int_{\frac{1}{n+1}}^{\pi}\right) \|(\phi(\cdot, t)) \sin^{\beta}(\cdot)\|_r |K_n^{EH}(t)| dt \\ &= O\left((n+1) \int_0^{\frac{1}{n+1}} t^{\beta} \omega(t) dt\right) + O\left(\frac{1}{(n+1)^2} \int_{\frac{1}{n+1}}^{\pi} t^{\beta-2} \omega(t) dt\right) \\ &= O\left((n+1) \omega\left(\frac{1}{n+1}\right) \int_0^{\frac{1}{n+1}} t^{\beta} dt\right) + O\left(\frac{1}{(n+1)^2} \int_{\frac{1}{n+1}}^{\pi} \frac{\omega(t)}{t^{2-\beta}} dt\right) \\ &= O\left(\frac{1}{(n+1)^{\beta}} \omega\left(\frac{1}{n+1}\right)\right) + O\left(\frac{1}{(n+1)^2} \int_{\frac{1}{n+1}}^{\pi} \frac{\omega(t)}{t^{2-\beta}} dt\right). \end{aligned} \quad (3.15)$$

Now from (3.14) and (3.15), we have

$$\begin{aligned}
\|(\mathcal{L}_n(\cdot)) \sin^\beta(\cdot)\|_r^v &= \|(\mathcal{L}_n(\cdot)) \sin^\beta(\cdot)\|_r \\
&\quad + \sup_{y \neq 0} \frac{\|(\mathcal{L}_n(\cdot + y) + \mathcal{L}_n(\cdot - y) - 2\mathcal{L}_n(\cdot)) \sin^\beta(\cdot)\|_r}{v(y)} \\
&= O\left(\frac{1}{(n+1)^\beta} w\left(\frac{1}{n+1}\right)\right) + O\left(\frac{1}{(n+1)^2} \int_{\frac{1}{n+1}}^\pi \frac{\omega(t)}{t^{2-\beta}} dt\right) \\
&\quad + O\left(\frac{1}{(n+1)^\beta} \frac{\omega\left(\frac{1}{n+1}\right)}{v\left(\frac{1}{n+1}\right)}\right) \\
&\quad + O\left(\frac{1}{(n+1)} \int_{\frac{1}{(n+1)^2}}^\pi \frac{t^{\beta-2}\omega(t)}{v(t)} dt\right) \\
&= \sum_{i=1}^4 O(J_i) \text{ (say)}. \tag{3.16}
\end{aligned}$$

Now we write  $J_1$  in terms of  $J_3$  and further  $J_2, J_3$  in terms of  $J_4$ . In view of monotonicity of  $v(t)$  for  $0 < t \leq \pi$ , we have

$$\omega(t) = \frac{\omega(t)}{v(t)} \cdot v(t) \leq v(\pi) \frac{\omega(t)}{v(t)} \cdot v(t) = O\left(\frac{\omega(t)}{v(t)}\right) \text{ for } 0 < t \leq \pi.$$

Therefore, we can write for  $t = (n+1)^{-1}$ .

$$J_1 = O(J_3). \tag{3.17}$$

Again by using monotonicity of  $v(t)$ ,

$$\begin{aligned}
J_2 &= \frac{1}{(n+1)^2} \int_{\frac{1}{n+1}}^\pi \frac{t^{\beta-2}\omega(t)}{v(t)} v(t) dt \\
&\leq \frac{1}{(n+1)^2} v(\pi) \int_{\frac{1}{n+1}}^\pi \frac{t^{\beta-2}\omega(t)}{v(t)} dt \\
&\leq \frac{1}{(n+1)^2} \int_{\frac{1}{n+1}}^\pi \frac{t^{\beta-2}\omega(t)}{v(t)} dt \\
&= O(J_4). \tag{3.18}
\end{aligned}$$

Now using  $\left(\frac{\omega(t)}{v(t)}\right)$  is positive and non-decreasing, we have

$$\begin{aligned}
J_4 &= \frac{1}{(n+1)} \int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-2} \omega(t)}{v(t)} dt \\
&\geq \frac{1}{(n+1)} \frac{\omega(\frac{1}{n+1})}{v(\frac{1}{n+1})} \int_{\frac{1}{n+1}}^{\pi} t^{\beta-2} dt \\
&\geq \frac{1}{(n+1)} \frac{\omega(\frac{1}{n+1})}{v(\frac{1}{n+1})} \frac{1}{(n+1)^{\beta-1}} \\
&\geq \frac{1}{(n+1)^{\beta}} \frac{\omega(\frac{1}{n+1})}{v(\frac{1}{n+1})}.
\end{aligned} \tag{3.19}$$

Hence, we have

$$J_3 = O(J_4). \tag{3.20}$$

Now combining (3.15) and (3.20), we get

$$\|(\mathcal{L}_n(\cdot)) \sin^{\beta}(\cdot)\|_r = O\left(\frac{1}{(n+1)} \int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-2} \omega(t)}{v(t)} dt\right). \tag{3.21}$$

Hence,

$$E_n(f) = \inf_n \|(\mathcal{L}_n(\cdot)) \sin^{\beta}(\cdot)\|_r^{(v)} = O\left(\frac{1}{(n+1)} \int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-2} \omega(t)}{v(t)} dt\right). \tag{3.22}$$

This completes the proof.  $\square$

**Theorem 3.6.** *Let  $\eta$  be a  $2\pi$  periodic function and Lebesgue integrable on  $[-\pi, \pi]$  and belonging to weighted Zygmund class  $W(Z_r^{(\omega)})$ ,  $r \geq 1$ . Then the degree of approximation of signal (function)  $\eta$ , using Euler-Hausdorff summability means of HL-series (2.3) is given by*

$$E_n(\eta) = \inf \|T_n^{EH} - \eta\|_r = O\left(\frac{1}{(n+1)} \frac{t^{\beta} \omega(\frac{1}{n+1})}{v(\frac{1}{n+1})}\right), \tag{3.23}$$

where  $\omega$  and  $v$  denotes the Zygmund moduli such that  $(\frac{\omega(t)}{tv(t)})$  is positive and decreasing.

*Proof.* Following the proof of Theorem 3.5, we have

$$E_n(f) = O\left(\frac{1}{(n+1)} \int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-2} \omega(t)}{v(t)} dt\right). \tag{3.24}$$

From our assumption that  $\left(\frac{\omega(t)}{tv(t)}\right)$  is positive and non-increasing with  $t$ , we have

$$\begin{aligned} E_n(f) &= O\left(\frac{1}{(n+1)}(n+1)\frac{\omega\left(\frac{1}{n+1}\right)}{v\left(\frac{1}{n+1}\right)}\int_{\frac{1}{n+1}}^{\pi} t^{\beta-1} dt\right) \\ &= O\left(\frac{\omega\left(\frac{1}{n+1}\right)}{v\left(\frac{1}{n+1}\right)}\left(\pi^\beta - \frac{1}{(n+1)^\beta}\right)\right). \end{aligned} \quad (3.25)$$

This completes the proof.  $\square$

#### 4. APPLICATIONS

Following corollaries can be obtained from Theorem 3.5.

**Corollary 4.1.** *If we replace Euler-Hausdorff mean by  $(E, 1)(C, 1)$  mean [11] in Theorem 3.5, then the degree of approximation of a function  $f \in W(Z_r^\omega)$  by  $(E, 1)(C, 1)$  mean of HL-series (2.3) is given by*

$$E_n(f) = O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-1}\omega(t)}{v(t)} dt\right). \quad (4.1)$$

**Corollary 4.2.** *If we replace Euler-Hausdorff mean by  $(E, q)(N, p_n, q_n)$  mean [8] in Theorem 3.5, then the degree of approximation of a function  $f \in W(Z_r^\omega)$  by  $(E, q)(N, p_n, q_n)$  mean of HL-series (2.3) is given by*

$$E_n(f) = O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-1}\omega(t)}{v(t)} dt\right). \quad (4.2)$$

**Corollary 4.3.** *If we replace Euler-Hausdorff mean by Hausdorff mean [5] in Theorem 3.5, then the degree of approximation of a function  $f \in W(Z_r^\omega)$  by Hausdorff mean of HL-series (2.3) is given by*

$$E_n(f) = O\left(\frac{1}{(n+1)}\int_{\frac{1}{n+1}}^{\pi} \frac{t^{\beta-1}\omega(t)}{v(t)} dt\right). \quad (4.3)$$

#### 5. CONCLUSION

There are various types of results exist in the literature concerning the degree of approximations of periodic functions belonging to different Zygmund classes and weighted Zygmund classes. The established theorem in this paper is an attempt to study the approximation of signals (functions) belonging to weighted Zygmund class via Euler-Hausdorff summability means for Hardy-Littlewood series, which generalizes several known results. Further, the result

can be extended for other functions belonging to weighted Zygmund class using Fourier series, conjugate Fourier series, derived Fourier series.

## REFERENCES

- [1] A.A. Das, S.K. Paikray, T. Pradhan and H. Dutta, *Approximation of signals in the weighted Zygmund class via Euler-Huasdorff product summability mean of Fourier series*, J. Indian Math. Soc., **87**(1–2) (2020), 22–36.
- [2] G.H. Hardy, *Divergent Series*, Oxford University Press, First Edition, 1949.
- [3] W.A. Hurwitz and L.L. Silverman, *On the consistency and equivalence of certain definitions of summability*, Trans. Amer. Math. Soc., **18** (1917), 1–20.
- [4] B.B. Jena, S.K. Paikray and M. Mursaleen, *On degree of approximation of Fourier series based on a certain class of product deferred summability means*, J. Inequal. Appl., **2023**(18) (2023), doi.org.10.1186/s13660-023-02927-z.
- [5] S. Lal and A. Mishra, *Euler-Hausdorff matrix summability operator and trigonometric approximation of the conjugate of a function belonging to the generalized Lipschitz class*, J. Inequal. Appl., **59** (2013), <https://doi.org/10.1186/1029-242X-2013-59>.
- [6] S. Lal and Shireen, *Best approximation of functions of generalized Zygmund class by Matrix-Euler summability mean of Fourier series*, Bull. Math. Anal. Appl., **5**(4) (2013), 1–13.
- [7] L. Leindler, *Strong approximation and generalized Zygmund class*, Acta Sci. Math., **43** (1981), 301–309.
- [8] M. Misra, P. Palo, B.P. Padhy, P. Samanta and U.K. Misra, *Approximation of Fourier series of a function of Lipschitz class by product means*, J. Adv. Math., **9**(4) (2014), 2475–2483.
- [9] F. Moricz, *Enlarged Lipschitz and Zygmund classes of functions and Fourier transforms*, East J. Approx., **16**(3) (2010), 259–271.
- [10] F. Moricz and J. Nemeth, *Generalized Zygmund classes of functions and strong approximation of Fourier series*, Acta. Sci. Math., **73** (2007), 637–647.
- [11] H.K. Nigam and K. Sharma, *On  $(E,1)(C,1)$  summability of Fourier series and its conjugate series*, Int. J. Pure Appl. Math., **82**(3) (2013), 365–375.
- [12] B.P. Padhy, P. Baliarsingh, L. Nayak and H. Dutta, *Approximation of Functions belonging to Zygmund Class Associated with Hardy-Littlewood Series using Riesz Mean*, Appl. Math. Inform. Sci., **16**(2) (2022), 243–248.
- [13] T. Pradhan, S.K. Paikray, A.A. Das and H. Dutta, *On approximation of signals in the generalized Zygmund class via  $(E,1)(\bar{N}, p_n)$  summability means of conjugate Fourier series*, Proyecciones J. Math., **38**(5) (2019), 981–998.
- [14] T. Pradhan, S.K. Paikray and U. Misra, *Approximation of signals belonging to generalized Lipschitz class using  $(\bar{N}, p_n, q_n)(E, s)$ -summability mean of Fourier series*, Cogent Math., **3**(1) (2016), 1–9.
- [15] T. Pradhan, S.K. Paikray and U.K. Misra, *Approximation of signals using generalized Zygmund class using  $(E,1)(\bar{N}, p_n)$  summability means of Fourier series*, Indian Soc. Indust. Appl. Math., **10**(1) (2019), 152–164.
- [16] M.V. Singh, M.L. Mittal and B.E. Rhoades, *Approximation of functions in the generalized Zygmund class using Hausdorff means*, J. Inequal. Appl., **101** (2017), DOI 10.1186/s13600-017-1361-8.
- [17] E.C. Titchmarsh, *The Theory of Functions*, Oxford University Press, 1939.
- [18] A. Zygmund, *Trigonometric series*, 2nd rev. ed., I, Cambridge Univ. Press, Cambridge, **51**, 1968.